



## Some regions *demand* polar coordinates

Some regions are best described in polar coordinates:



Parts of circles



Parts of annuli


These are called **polar rectangles** because they have the form

$$\left\{ \begin{array}{l} a \leq r \leq b \\ \alpha \leq \theta \leq \beta \end{array} \right\} \text{ for constants } \left\{ \begin{array}{l} a, b \\ \alpha, \beta \end{array} \right\}$$

# Polar coordinates

**Goal:** Convert a double integral involving  $x$ 's and  $y$ 's into a double integral involving  $r$ 's and  $\theta$ 's.

**Important:** Using “polar slices” introduces a complication.



In this picture,  $dA$  is **not**  $dr d\theta$ .

The radial component is \_\_\_\_\_ and the circular component is \_\_\_\_\_.

This means  $dA =$  \_\_\_\_\_. (How to remember?)

**Consequence:** To calculate  $\iint_D f(x, y) dA$ , when  $D$  is best described in polar coordinates, calculate


$$\iint_D f(r \cos \theta, r \sin \theta) r dr d\theta$$

## Polar coordinate example

**Example.** Find the area inside  
 $r = 4 \cos \theta$  from  $\theta = \frac{\pi}{4}$  to  $\frac{\pi}{2}$ .

This region is defined as  $\frac{\pi}{4} \leq \theta \leq \frac{\pi}{2}$   
 and \_\_\_\_\_  $\leq r \leq$  \_\_\_\_\_.

We use  $A = \iint dA$ :



★ Along the way, we had  $\int_{\alpha}^{\beta} \frac{r^2}{2} d\theta \dots\dots\dots$

## Application: Density (p. 692/723)

**Example.** The density of a point on a semicircular lamina of radius  $a$  is proportional to the distance from the center of the circle. Find the mass of the lamina.

**Solution. Draw a picture!**

**Setup:** Let the lamina be the upper half of the circle  $x^2 + y^2 = a^2$ , which is a polar rectangle:

The density function can be written:

$$\rho(x, y) = K\sqrt{x^2 + y^2}$$

The total mass is  $m = \iint_D \rho(x, y) dA$

$$\dots = \int_{\theta=0}^{\theta=\pi} K \frac{a^3}{3} d\theta = \left[ K \frac{a^3}{3} \theta \right]_{\theta=0}^{\theta=\pi} = \frac{\pi K a^3}{3}.$$

## Changing from $(x, y)$ to $(r, \theta)$

**Example.** Calculate  $\int_0^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} (x^3 + xy^2) dy dx$  (hint for polar!)

**Solution.** Draw a picture!

Notice that this is the polar rectangle  
 $0 \leq r \leq 3$  and  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ .

Rewrite the integral as

$$\begin{aligned} & \iint_D (r^3 \cos^3 \theta + r^3 \cos \theta \sin^2 \theta) r dr d\theta \\ &= \iint_D r^4 \cos \theta (\cos^2 \theta + \sin^2 \theta) dr d\theta = \iint_D r^4 \cos \theta dr d\theta \\ &= \left( \int_{r=0}^{r=3} r^4 dr \right) \cdot \left( \int_{\theta=-\pi/2}^{\theta=\pi/2} \cos \theta d\theta \right) \\ &= \left[ \frac{r^5}{5} \right]_{r=0}^{r=3} \cdot [\sin \theta]_{\theta=-\pi/2}^{\theta=\pi/2} = \left( \frac{3^5}{5} - 0 \right) \cdot (1 - (-1)) = \frac{2 \cdot 3^5}{5} \end{aligned}$$

## Volume example

**Example.** Find the volume of the solid under the paraboloid  $z = x^2 + y^2$ , above the  $xy$ -plane, and inside the cylinder  $(x - 1)^2 + y^2 = 1$ .

**Plan of attack: Draw a picture!**

- ▶ Notice circley thingees. Think: Possible Polar Problem.
- ▶ Convert the given information to polar equations using

$$(x, y) = (r \cos \theta, r \sin \theta):$$

$$(x-1)^2 + y^2 = 1 \rightsquigarrow (r \cos \theta - 1)^2 + (r \sin \theta)^2 = 1 \rightsquigarrow r = 2 \cos \theta.$$

$$(x^2 + y^2) \rightsquigarrow r^2.$$

- ▶ Set up the polar integral.

$$\text{So } \iint_D (x^2 + y^2) dA = \iint_D r^2 r dr d\theta.$$

- ▶ Integrate.....