

# Functions

## Single-variable functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f : x \mapsto f(x)$$

$f$  takes in a real number  $x$   
outputs a real number  $f(x)$

## Vector functions

$$\vec{r} : \mathbb{R} \rightarrow \mathbb{R}^3 \quad (\text{or } \mathbb{R}^2 \text{ or } \mathbb{R}^n)$$

$$\vec{r} : t \mapsto \langle f(t), g(t), h(t) \rangle$$

$\vec{r}$  takes in a real number  $t$   
outputs a vector  $\langle f(t), g(t), h(t) \rangle$

## Limits and Helices

The **limit** of a vector function  $\vec{r}$  is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

A vector-valued function  $\vec{r}(t)$  is continuous at  $a$  if \_\_\_\_\_.

**Example.** Sketch the curve given by  $\vec{r}(t) = \cos t \vec{i} + \sin t \vec{j} + t \vec{k}$ .

The  $x$  and  $y$  components \_\_\_\_\_ while the  $z$  component \_\_\_\_\_.

Plug in some values of  $t$

# Intersectionnnnnnnnnnnnn

**Example.** Find a vector function that is the intersection of the cylinder  $x^2 + z^2 = 1$  and the plane  $y + z = 2$ .

**Strategems:** Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as  $t$ .
- ▶ When the curve is closed, this is not possible—work first in 2D.

**Answer:** Use the fact that we are on the cylinder. (Eqn 1) Project onto the  $xz$ -plane and start the parametrization there:

$$x(t) = \quad z(t) = \quad \underline{\quad} \leq t \leq \underline{\quad}.$$

Use (Eqn 2) to find the  $y$ -coordinate:

So  $\vec{r} =$

## Derivatives and Derivative-derivative definitions

Define  $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$ .

The **derivative**  $\vec{r}'(t)$  of a vector-valued function is a vector in the direction tangent to the curve  $\vec{r}(t)$ .

- ▶ Standardize. The **unit tangent vector**  $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$ .
- ▶ We can take multiple derivatives  $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$
- ▶ A function is **smooth** on an interval  $I$  if
  - ▶  $\vec{r}'(t)$  is continuous on  $I$
  - ▶ and  $\vec{r}'(t) \neq \vec{0}$ , except possibly at the endpoints of  $I$

We can integrate too.  $\int_a^b \vec{r}(t) dt = \langle \int_a^b f(t) dt, \int_a^b g(t) dt, \int_a^b h(t) dt \rangle$

Remember: Indefinite integrals have a (vector) constant of integration.

**Example.**  $\int \langle 2 \cos t, \sin t, 2t \rangle dt = \langle 2 \sin t, -\cos t, t^2 \rangle + \vec{C}$ .

# Derivatives

**Example.** Find the equation of the tangent line to the helix  $\vec{r}(t) = \langle 2 \cos t, \sin t, t \rangle$  at the point  $P = (0, 1, \frac{\pi}{2})$ .

## Game plan:

1. Find  $t^*$  for which the curve goes through the point  $P$ .
2. Find the tangent vector  $\vec{r}'(t)$ , plug in  $t = t^*$ .
3. Write the equation of the line.

# Derivatives rule

- ▶  $\frac{d}{dt} (\vec{\mathbf{r}}(t) + \vec{\mathbf{s}}(t)) = \vec{\mathbf{r}}'(t) + \vec{\mathbf{s}}'(t)$
- ▶  $\frac{d}{dt} (c \vec{\mathbf{r}}(t)) = c \vec{\mathbf{r}}'(t)$
- ▶  $\frac{d}{dt} (f(t) \vec{\mathbf{r}}(t)) = f'(t) \vec{\mathbf{r}}(t) + f(t) \vec{\mathbf{r}}'(t)$
- ▶  $\frac{d}{dt} (\vec{\mathbf{r}}(t) \cdot \vec{\mathbf{s}}(t)) = \vec{\mathbf{r}}'(t) \cdot \vec{\mathbf{s}}(t) + \vec{\mathbf{r}}(t) \cdot \vec{\mathbf{s}}'(t)$
- ▶  $\frac{d}{dt} (\vec{\mathbf{r}}(t) \times \vec{\mathbf{s}}(t)) = \vec{\mathbf{r}}'(t) \times \vec{\mathbf{s}}(t) + \vec{\mathbf{r}}(t) \times \vec{\mathbf{s}}'(t)$
- ▶  $\frac{d}{dt} (\vec{\mathbf{r}}(f(t))) = f'(t) \vec{\mathbf{r}}'(f(t))$

## Motion in space

If  $\vec{r}(t)$  is the vector position of a particle, then

- ▶  $\vec{r}'(t) = \vec{v}(t)$  is the vector velocity of the particle.
- ▶  $|\vec{r}'(t)| = |\vec{v}(t)| = \text{speed of the particle.}$
- ▶  $\vec{r}''(t) = \vec{a}(t)$  is the vector acceleration of the particle.

We can use  $\vec{a}(t)$  to find the force that an object exerts:  $\vec{F}(t) = m\vec{a}(t)$

**Example.** Suppose that a mass of 40 kg starts with init. pos'n  $\langle 1, 0, 0 \rangle$ , initial velocity  $\langle 1, -1, 1 \rangle$  and has acceleration  $\vec{a}(t) = \langle 4t, 6t, 1 \rangle$ .

- (a) Find the position and velocity of the particle as a function of  $t$ .
- (b) Determine the force that the particle exerts at time  $t = 2$ .

**Example.** Show that if a particle moves with constant speed, then the velocity and acceleration vectors are orthogonal.