

Functions

Single-variable functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f : x \mapsto f(x)$$

f takes in a real number x
outputs a real number $f(x)$

Functions

Single-variable functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f : x \mapsto f(x)$$

f takes in a real number x
outputs a real number $f(x)$

Vector functions

$$\vec{r} : \mathbb{R} \rightarrow \mathbb{R}^3 \quad (\text{or } \mathbb{R}^2 \text{ or } \mathbb{R}^n)$$

$$\vec{r} : t \mapsto \langle f(t), g(t), h(t) \rangle$$

\vec{r} takes in a real number t
outputs a vector $\langle f(t), g(t), h(t) \rangle$

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

A vector-valued function $\vec{r}(t)$ is continuous at a if _____.

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

A vector-valued function $\vec{r}(t)$ is continuous at a if _____.

Example. Sketch the curve given by $\vec{r}(t) = \cos t \vec{i} + \sin t \vec{j} + t \vec{k}$.

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

A vector-valued function $\vec{r}(t)$ is continuous at a if _____.

Example. Sketch the curve given by $\vec{r}(t) = \cos t \vec{i} + \sin t \vec{j} + t \vec{k}$.
The x and y components _____ while the z component _____.

Limits and Helices

The **limit** of a vector function \vec{r} is defined by taking the limits of its component functions (as long as each of these exists...)

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

A vector-valued function $\vec{r}(t)$ is continuous at a if _____.

Example. Sketch the curve given by $\vec{r}(t) = \cos t \vec{i} + \sin t \vec{j} + t \vec{k}$.
The x and y components _____ while the z component _____.
Plug in some values of t

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization...

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as t .

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as t .
- ▶ When the curve is closed, this is not possible—work first in 2D.

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as t .
- ▶ When the curve is closed, this is not possible—work first in 2D.

Answer: Use the fact that we are on the cylinder. (Eqn 1) Project onto the xz -plane and start the parametrization there:

$$x(t) = \qquad z(t) = \qquad \text{____} \leq t \leq \text{____}.$$

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as t .
- ▶ When the curve is closed, this is not possible—work first in 2D.

Answer: Use the fact that we are on the cylinder. (Eqn 1) Project onto the xz -plane and start the parametrization there:

$$x(t) = \quad \quad \quad z(t) = \quad \quad \quad \underline{\quad} \leq t \leq \underline{\quad}.$$

Use (Eqn 2) to find the y -coordinate:

Intersectionnnnnnnnnnn

Example. Find a vector function that is the intersection of the cylinder $x^2 + z^2 = 1$ and the plane $y + z = 2$.

Strategems: Find a parametrization... What parameter to use?

- ▶ If a curve is oriented in one direction, use that variable as t .
- ▶ When the curve is closed, this is not possible—work first in 2D.

Answer: Use the fact that we are on the cylinder. (Eqn 1) Project onto the xz -plane and start the parametrization there:

$$x(t) = \quad \quad \quad z(t) = \quad \quad \quad \text{______} \leq t \leq \text{______}.$$

Use (Eqn 2) to find the y -coordinate:

So $\vec{r} =$

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.
- ▶ We can take multiple derivatives $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.
- ▶ We can take multiple derivatives $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$
- ▶ A function is **smooth** on an interval I if
 - ▶ $\vec{r}'(t)$ is continuous on I
 - ▶ and $\vec{r}'(t) \neq \vec{0}$, except possibly at the endpoints of I

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.
- ▶ We can take multiple derivatives $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$
- ▶ A function is **smooth** on an interval I if
 - ▶ $\vec{r}'(t)$ is continuous on I
 - ▶ and $\vec{r}'(t) \neq \vec{0}$, except possibly at the endpoints of I

We can integrate too. $\int_a^b \vec{r}(t) dt = \langle \int_a^b f(t) dt, \int_a^b g(t) dt, \int_a^b h(t) dt \rangle$

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.
- ▶ We can take multiple derivatives $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$
- ▶ A function is **smooth** on an interval I if
 - ▶ $\vec{r}'(t)$ is continuous on I
 - ▶ and $\vec{r}'(t) \neq \vec{0}$, except possibly at the endpoints of I

We can integrate too. $\int_a^b \vec{r}(t) dt = \langle \int_a^b f(t) dt, \int_a^b g(t) dt, \int_a^b h(t) dt \rangle$

Remember: Indefinite integrals have a (vector) constant of integration.

Derivatives and Derivative-derivative definitions

Define $\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} = \langle f'(t), g'(t), h'(t) \rangle$.

The **derivative** $\vec{r}'(t)$ of a vector-valued function is a vector in the direction tangent to the curve $\vec{r}(t)$.

- ▶ Standardize. The **unit tangent vector** $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$.
- ▶ We can take multiple derivatives $\vec{r}''(t) = \frac{d}{dt}(\vec{r}'(t))$
- ▶ A function is **smooth** on an interval I if
 - ▶ $\vec{r}'(t)$ is continuous on I
 - ▶ and $\vec{r}'(t) \neq \vec{0}$, except possibly at the endpoints of I

We can integrate too. $\int_a^b \vec{r}(t) dt = \langle \int_a^b f(t) dt, \int_a^b g(t) dt, \int_a^b h(t) dt \rangle$

Remember: Indefinite integrals have a (vector) constant of integration.

Example. $\int \langle 2 \cos t, \sin t, 2t \rangle dt = \langle 2 \sin t, -\cos t, t^2 \rangle + \vec{C}$.

Derivatives

Example. Find the equation of the tangent line to the helix $\vec{r}(t) = \langle 2 \cos t, \sin t, t \rangle$ at the point $P = (0, 1, \frac{\pi}{2})$.

Derivatives

Example. Find the equation of the tangent line to the helix $\vec{r}(t) = \langle 2 \cos t, \sin t, t \rangle$ at the point $P = (0, 1, \frac{\pi}{2})$.

Game plan:

1. Find t^* for which the curve goes through the point P .
2. Find the tangent vector $\vec{r}'(t)$, plug in $t = t^*$.
3. Write the equation of the line.

Derivatives rule

- ▶ $\frac{d}{dt}(\vec{r}(t) + \vec{s}(t)) = \vec{r}'(t) + \vec{s}'(t)$
- ▶ $\frac{d}{dt}(c \vec{r}(t)) = c \vec{r}'(t)$
- ▶ $\frac{d}{dt}(f(t) \vec{r}(t)) = f'(t) \vec{r}(t) + f(t) \vec{r}'(t)$
- ▶ $\frac{d}{dt}(\vec{r}(t) \cdot \vec{s}(t)) = \vec{r}'(t) \cdot \vec{s}(t) + \vec{r}(t) \cdot \vec{s}'(t)$
- ▶ $\frac{d}{dt}(\vec{r}(t) \times \vec{s}(t)) = \vec{r}'(t) \times \vec{s}(t) + \vec{r}(t) \times \vec{s}'(t)$
- ▶ $\frac{d}{dt}(\vec{r}(f(t))) = f'(t) \vec{r}'(f(t))$

Motion in space

If $\vec{r}(t)$ is the vector position of a particle, then

- ▶ $\vec{r}'(t) = \vec{v}(t)$ is the vector velocity of the particle.
- ▶ $|\vec{r}'(t)| = |\vec{v}(t)| = \text{speed of the particle.}$
- ▶ $\vec{r}''(t) = \vec{a}(t)$ is the vector acceleration of the particle.

Motion in space

If $\vec{r}(t)$ is the vector position of a particle, then

- ▶ $\vec{r}'(t) = \vec{v}(t)$ is the vector velocity of the particle.
- ▶ $|\vec{r}'(t)| = |\vec{v}(t)| = \text{speed of the particle.}$
- ▶ $\vec{r}''(t) = \vec{a}(t)$ is the vector acceleration of the particle.

We can use $\vec{a}(t)$ to find the force that an object exerts: $\vec{F}(t) = m\vec{a}(t)$

Motion in space

If $\vec{r}(t)$ is the vector position of a particle, then

- ▶ $\vec{r}'(t) = \vec{v}(t)$ is the vector velocity of the particle.
- ▶ $|\vec{r}'(t)| = |\vec{v}(t)| = \text{speed of the particle.}$
- ▶ $\vec{r}''(t) = \vec{a}(t)$ is the vector acceleration of the particle.

We can use $\vec{a}(t)$ to find the force that an object exerts: $\vec{F}(t) = m\vec{a}(t)$

Example. Suppose that a mass of 40 kg starts with init. pos'n $\langle 1, 0, 0 \rangle$, initial velocity $\langle 1, -1, 1 \rangle$ and has acceleration $\vec{a}(t) = \langle 4t, 6t, 1 \rangle$.

Motion in space

If $\vec{r}(t)$ is the vector position of a particle, then

- ▶ $\vec{r}'(t) = \vec{v}(t)$ is the vector velocity of the particle.
- ▶ $|\vec{r}'(t)| = |\vec{v}(t)| = \text{speed of the particle.}$
- ▶ $\vec{r}''(t) = \vec{a}(t)$ is the vector acceleration of the particle.

We can use $\vec{a}(t)$ to find the force that an object exerts: $\vec{F}(t) = m\vec{a}(t)$

Example. Suppose that a mass of 40 kg starts with init. pos'n $\langle 1, 0, 0 \rangle$, initial velocity $\langle 1, -1, 1 \rangle$ and has acceleration $\vec{a}(t) = \langle 4t, 6t, 1 \rangle$.

- Find the position and velocity of the particle as a function of t .
- Determine the force that the particle exerts at time $t = 2$.

Motion in space

If $\vec{r}(t)$ is the vector position of a particle, then

- ▶ $\vec{r}'(t) = \vec{v}(t)$ is the vector velocity of the particle.
- ▶ $|\vec{r}'(t)| = |\vec{v}(t)| =$ speed of the particle.
- ▶ $\vec{r}''(t) = \vec{a}(t)$ is the vector acceleration of the particle.

We can use $\vec{a}(t)$ to find the force that an object exerts: $\vec{F}(t) = m\vec{a}(t)$

Example. Suppose that a mass of 40 kg starts with init. pos'n $\langle 1, 0, 0 \rangle$, initial velocity $\langle 1, -1, 1 \rangle$ and has acceleration $\vec{a}(t) = \langle 4t, 6t, 1 \rangle$.

- Find the position and velocity of the particle as a function of t .
- Determine the force that the particle exerts at time $t = 2$.

Example. Show that if a particle moves with constant speed, then the velocity and acceleration vectors are orthogonal.