

Chain Rule

Function of one variable

Suppose $y = f(x)$ and $x = g(t)$.

That is, $y = f(g(t))$.

The chain rule gives:

$$\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt}$$

$$\frac{dy}{dt} = f'(g(t)) \cdot g'(t)$$

Key idea:

You must add contributions from all dependencies.

Function of several variables

Suppose $z = f(x, y)$ and $\begin{cases} x = g(t) \\ y = h(t) \end{cases}$

So $z = f(g(t), h(t))$.

The chain rule gives

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \cdot \frac{dy}{dt}$$

$$\frac{dz}{dt} = f_x(g(t), h(t)) \cdot g'(t) + f_y(g(t), h(t)) \cdot h'(t)$$

Example. Let $z = x^2y + 3xy^2$, where $x = \sin 2t$, $y = \cos t$. Find $\frac{dz}{dt}$, $z'(0)$.

Answer:

More Chains

All dependencies

Alternatively, we might have $z = f(x, y)$

and $x = g(s, t)$, $y = h(s, t)$.

Then $\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s}$.

Example. Consider $u = x^4 y + y^2 z^3$ where $x = rse^t$, $y = rs^2 e^{-t}$, $z = r^2 s(\sin t)$. Find $\frac{\partial u}{\partial s}$.

In full generality: If u is a function of x_1, x_2, \dots, x_n and each x_j is a function of t_1, t_2, \dots, t_m , then

$$\frac{\partial u}{\partial t_i} = \frac{\partial u}{\partial x_1} \cdot \frac{\partial x_1}{\partial t_i} + \frac{\partial u}{\partial x_2} \cdot \frac{\partial x_2}{\partial t_i} + \dots + \frac{\partial u}{\partial x_n} \cdot \frac{\partial x_n}{\partial t_i}.$$

Implicit differentiation

- Simplify implicit differentiation calculations!

Involving two variables

Consider $F(x, y) = C$ like

Implicitly y is a function of x :

$$F(x, f(x)) = C$$

Differentiating w.r.t. x :

$$F(x, y) = C$$

$$\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} = 0$$

Solving for $\frac{dy}{dx}$ gives

$$\frac{dy}{dx} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial y}}$$

Involving three variables

Consider $F(x, y, z) = C$

Implicitly z is a function of x and y :

$$F(x, y, f(x, y)) = C$$

Differentiating w.r.t. x :

$$F(x, y, z) = C$$

$$\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} + \frac{\partial F}{\partial z} \frac{dz}{dx} = 0$$

Solving for $\frac{dz}{dx}$ gives

$$\frac{dz}{dx} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}}$$

Example. Find $\frac{\partial z}{\partial x}$ if $x^3 + y^3 + z^3 + 6xyz = 1$