

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a}^{x=b} y \, dx$$

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a}^{x=b} y \, dx = \int_{t=\alpha}^{t=\beta} g(t) f'(t) \, dt$$

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
 (Here,  $r$  is a constant and  $\theta$  is the parameter.)

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
 (Here,  $r$  is a constant and  $\theta$  is the parameter.)

Plot it to see the shape.

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
 (Here,  $r$  is a constant and  $\theta$  is the parameter.)

Plot it to see the shape. One arch has range  $\underline{\quad} \leq \theta \leq \underline{\quad}$ .

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
 (Here,  $r$  is a constant and  $\theta$  is the parameter.)

Plot it to see the shape. One arch has range  $\_\_\_ \leq \theta \leq \_\_\_\_\_$ .

$$A = \int y \, dx = \int r(1 - \cos \theta) r(1 - \cos \theta) \, d\theta$$

## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
(Here,  $r$  is a constant and  $\theta$  is the parameter.)

Plot it to see the shape. One arch has range  $\underline{\quad} \leq \theta \leq \underline{\quad}$ .

$$\begin{aligned} A &= \int y \, dx = \int r(1 - \cos \theta) r(1 - \cos \theta) \, d\theta \\ &= r^2 \int (1 - 2 \cos \theta + \cos^2 \theta) \, d\theta \end{aligned}$$



## Area under a parametric curve

Given  $y = f(x)$ , the area under the curve from  $x = a$  to  $x = b$  is

$$\text{Area} = \int_{x=a \leftarrow \text{left endpoint}}^{x=b \leftarrow \text{right endpoint}} y \, dx = \int_{t=\alpha \leftarrow \text{left endpoint}}^{t=\beta \leftarrow \text{right endpoint}} g(t) f'(t) \, dt$$

**Example.** Find the area under one arch of the cycloid  $\begin{cases} x = r(\theta - \sin \theta) \\ y = r(1 - \cos \theta) \end{cases}$   
(Here,  $r$  is a constant and  $\theta$  is the parameter.)

Plot it to see the shape. One arch has range  $\_\_\_ \leq \theta \leq \_\_\_\_\_$ .

$$\begin{aligned} A &= \int y \, dx = \int r(1 - \cos \theta) r(1 - \cos \theta) \, d\theta \\ &= r^2 \int (1 - 2 \cos \theta + \cos^2 \theta) \, d\theta \end{aligned}$$

$$= 3\pi r^2$$

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

### Polar land

How much area is swept out by a little slice?

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

$$A = \int_{\theta=a}^{\theta=b} \frac{1}{2} [f(\theta)]^2 d\theta = \int_a^b \frac{1}{2} r^2 d\theta$$

### Polar land

How much area is swept out by a little slice?

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

$$A = \int_{\theta=a}^{\theta=b} \frac{1}{2} [f(\theta)]^2 \, d\theta = \int_a^b \frac{1}{2} r^2 \, d\theta$$

### Polar land

How much area is swept out by a little slice?

**Example.** What is the area inside one loop of the four-leaved rose  $r = \cos 2\theta$ ?

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

$$A = \int_{\theta=a}^{\theta=b} \frac{1}{2} [f(\theta)]^2 \, d\theta = \int_a^b \frac{1}{2} r^2 \, d\theta$$

### Polar land

How much area is swept out by a little slice?

**Example.** What is the area inside one loop of the four-leaved rose  $r = \cos 2\theta$ ?

- ▶ What are the bounds on  $\theta$ ?



## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

$$A = \int_{\theta=a}^{\theta=b} \frac{1}{2} [f(\theta)]^2 d\theta = \int_a^b \frac{1}{2} r^2 d\theta$$

### Polar land

How much area is swept out by a little slice?

**Example.** What is the area inside one loop of the four-leaved rose  $r = \cos 2\theta$ ?

- ▶ What are the bounds on  $\theta$ ?
- ▶ For which  $\theta$  does the curve pass through the origin?

## Area inside a polar curve

For cartesian functions  $y = f(x)$ , calculate area as  $A = \int dA = \int y \, dx$ .

What about the area “*inside a curve*” best described as a polar function  $r = f(\theta)$ ?

- ▶ We still use  $A = \int dA$ .
- ▶ But the formula for  $dA$  is different.

$$A = \int_{\theta=a}^{\theta=b} \frac{1}{2} [f(\theta)]^2 d\theta = \int_a^b \frac{1}{2} r^2 d\theta$$

### Polar land

How much area is swept out by a little slice?

**Example.** What is the area inside one loop of the four-leaved rose  $r = \cos 2\theta$ ?

- ▶ What are the bounds on  $\theta$ ?
- ▶ For which  $\theta$  does the curve pass through the origin?

$$A = \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \frac{1}{2} [\underline{\quad}]^2 d\theta$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

Now calculate:  $\int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned} \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\ &= \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta \end{aligned}$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned}
 \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 8 \sin^2 \theta - 2 \sin \theta - 1 \right] d\theta
 \end{aligned}$$



## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned}
 \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 8 \sin^2 \theta - 2 \sin \theta - 1 \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 4 - 4 \cos 2\theta - 2 \sin \theta - 1 \right] d\theta
 \end{aligned}$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned}
 \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 8 \sin^2 \theta - 2 \sin \theta - 1 \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 4 - 4 \cos 2\theta - 2 \sin \theta - 1 \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 3 - 4 \cos 2\theta - 2 \sin \theta \right] d\theta
 \end{aligned}$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned}
 \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} [9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1)] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} [8 \sin^2 \theta - 2 \sin \theta - 1] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} [4 - 4 \cos 2\theta - 2 \sin \theta - 1] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} [3 - 4 \cos 2\theta - 2 \sin \theta] d\theta \\
 = & \left[ \frac{3\theta}{2} - \sin 2\theta + \cos \theta \right]_{\pi/6}^{5\pi/6}
 \end{aligned}$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

$$\begin{aligned}
 \text{Now calculate: } & \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 8 \sin^2 \theta - 2 \sin \theta - 1 \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 4 - 4 \cos 2\theta - 2 \sin \theta - 1 \right] d\theta \\
 = & \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 3 - 4 \cos 2\theta - 2 \sin \theta \right] d\theta \\
 = & \left[ \frac{3\theta}{2} - \sin 2\theta + \cos \theta \right]_{\pi/6}^{5\pi/6} \\
 = & \frac{3}{2} \left( \frac{5\pi}{6} - \frac{\pi}{6} \right) - \left( \sin \frac{10\pi}{6} - \sin \frac{2\pi}{6} \right) + \left( \cos \frac{5\pi}{6} - \cos \frac{\pi}{6} \right) = \dots
 \end{aligned}$$

## Inside yet Outside

**Example.** Calculate the area inside the curve  $r = 3 \sin \theta$  and outside the curve  $r = \sin \theta + 1$ .

**First:** Draw a picture!

- ▶ What are these curves?
- ▶ Where do they intersect?

Now calculate:  $\int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ \overset{\text{outside}}{\left(\frac{1}{2}(3 \sin \theta)^2\right)} - \overset{\text{inside}}{\left(\frac{1}{2}(\sin \theta + 1)^2\right)} \right] d\theta$

$$= \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 9 \sin^2 \theta - (\sin^2 \theta + 2 \sin \theta + 1) \right] d\theta$$

$$= \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 8 \sin^2 \theta - 2 \sin \theta - 1 \right] d\theta$$

$$= \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 4 - 4 \cos 2\theta - 2 \sin \theta - 1 \right] d\theta$$

$$= \frac{1}{2} \int_{\theta=\underline{\quad}}^{\theta=\underline{\quad}} \left[ 3 - 4 \cos 2\theta - 2 \sin \theta \right] d\theta$$

$$= \left[ \frac{3\theta}{2} - \sin 2\theta + \cos \theta \right]_{\pi/6}^{5\pi/6}$$

$$= \frac{3}{2} \left( \frac{5\pi}{6} - \frac{\pi}{6} \right) - \left( \sin \frac{10\pi}{6} - \sin \frac{2\pi}{6} \right) + \left( \cos \frac{5\pi}{6} - \cos \frac{\pi}{6} \right) = \dots$$

$$= \pi + \sqrt{3} - \sqrt{3} = \pi$$

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .



## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

$$L = \int_{t=\alpha}^{t=\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

$$L = \int_{t=\alpha}^{t=\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

**Example.** Find the arc length for the parametric curve  $\begin{cases} x = \sin 2t \\ y = \cos 2t \end{cases}$  for  $0 \leq t \leq 2\pi$ .

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

$$L = \int_{t=\alpha}^{t=\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

**Example.** Find the arc length for the parametric curve  $\begin{cases} x = \sin 2t \\ y = \cos 2t \end{cases}$  for  $0 \leq t \leq 2\pi$ . **What do we expect?**

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

$$L = \int_{t=\alpha}^{t=\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

**Example.** Find the arc length for the parametric curve  $\begin{cases} x = \sin 2t \\ y = \cos 2t \end{cases}$   
for  $0 \leq t \leq 2\pi$ . **What do we expect? What is this curve?**

## Arc length of a parametric curve

To find the arc length of a parametric curve, think  $L = \int dL$ .

How much arc length  $dL$  does the curve traverse in one time unit  $dt$ ?

Draw a right triangle:

To see  $dL = \sqrt{dx^2 + dy^2}$ .

How are each related to  $dt$ ?

$$dL = \sqrt{\left[\frac{dx}{dt} \cdot dt\right]^2 + \left[\frac{dy}{dt} \cdot dt\right]^2}$$

$$L = \int_{t=\alpha}^{t=\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

**Example.** Find the arc length for the parametric curve  $\begin{cases} x = \sin 2t \\ y = \cos 2t \end{cases}$  for  $0 \leq t \leq 2\pi$ . **What do we expect? What is this curve?**

$$\int_{t=0}^{t=2\pi} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt =$$

## Arc length of a polar curve

For a polar curve, calculate  $dL$  in terms of  $dr$  and  $d\theta$ :

The right triangle shows:

$$dL = \sqrt{(r d\theta)^2 + (dr)^2}$$

Take  $\theta$  as the parameter to get



## Arc length of a polar curve

For a polar curve, calculate  $dL$  in terms of  $dr$  and  $d\theta$ :

The right triangle shows:

$$dL = \sqrt{(r d\theta)^2 + (dr)^2}$$

Take  $\theta$  as the parameter to get

$$dL = \sqrt{[r d\theta]^2 + \left[\frac{dr}{d\theta} d\theta\right]^2}$$

## Arc length of a polar curve

For a polar curve, calculate  $dL$  in terms of  $dr$  and  $d\theta$ :

The right triangle shows:

$$dL = \sqrt{(r d\theta)^2 + (dr)^2}$$

Take  $\theta$  as the parameter to get

$$dL = \sqrt{[r d\theta]^2 + \left[\frac{dr}{d\theta} d\theta\right]^2} \text{ so } L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

## Arc length of a polar curve

For a polar curve, calculate  $dL$  in terms of  $dr$  and  $d\theta$ :

The right triangle shows:

$$dL = \sqrt{(r d\theta)^2 + (dr)^2}$$

Take  $\theta$  as the parameter to get

$$dL = \sqrt{[r d\theta]^2 + \left[\frac{dr}{d\theta} d\theta\right]^2} \text{ so } L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Alternatively, use the parametrization  $\begin{cases} x = f(\theta) \cos \theta \\ y = f(\theta) \sin \theta \end{cases}$ .

► Then 
$$L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{\left[\frac{d}{d\theta}(f(\theta) \cos \theta)\right]^2 + \left[\frac{d}{d\theta}(f(\theta) \sin \theta)\right]^2} d\theta$$

## Arc length of a polar curve

For a polar curve, calculate  $dL$  in terms of  $dr$  and  $d\theta$ :

The right triangle shows:

$$dL = \sqrt{(r d\theta)^2 + (dr)^2}$$

Take  $\theta$  as the parameter to get

$$dL = \sqrt{[r d\theta]^2 + \left[\frac{dr}{d\theta} d\theta\right]^2} \text{ so } L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Alternatively, use the parametrization  $\begin{cases} x = f(\theta) \cos \theta \\ y = f(\theta) \sin \theta \end{cases}$ .

► Then 
$$L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{\left[\frac{d}{d\theta}(f(\theta) \cos \theta)\right]^2 + \left[\frac{d}{d\theta}(f(\theta) \sin \theta)\right]^2} d\theta$$

... take derivatives & do the algebra ...

$$L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{[f(\theta)]^2 + \left[\frac{d}{d\theta}(f(\theta))\right]^2} d\theta$$