

Counting words

Definition: A **list** or **word** is an ordered sequence of objects.

Definition: A **k -list** or **k -word** is a list of length k .


- ▶ A **list** or **word** is always ordered and a **set** is always unordered.

Question: How many lists have three entries where

- ▶ The first two entries can be either A or B .
- ▶ The last entry is either 5 or 6.

Answer: We can solve this using a tree diagram:

Alternatively: Notice two *independent* choices for each character. Multiply $2 \cdot 2 \cdot 2 = 8$.



The Product Principle

This illustrates:

The product principle: When counting lists (l_1, l_2, \dots, l_k) ,


IF there are c_1 choices for entry l_1 , each leading to a different list,

AND IF there are c_i choices for entry l_i ,

no matter the choices made for l_1 through l_{i-1} ,
each leading to a different list

THEN there are $c_1 c_2 \cdots c_k$ such lists.

Caution: The product principle seems simple, but we must be careful when we use it.



Lists WITH repetition

Q1. How many 8-character passwords are there using $A-Z$, $a-z$, $0-9$?

Answer: Creating a word of length 8, with ____ choices for each character. Therefore, the number of 8-character passwords is ____.
(=218,340,105,584,896)

In general, the number of words of length k that can be made from an alphabet of length n and where repetition is allowed is n^k

Application: Counting Subsets

Example. How many subsets of a set $S = \{s_1, s_2, \dots, s_n\}$ are there?

Strategy: “Try small problems, see a pattern.”

- ▶ $n = 0$: $S = \emptyset \rightsquigarrow \{\emptyset\}$, size 1.
- ▶ $n = 1$: $S = \{s_1\} \rightsquigarrow \{\emptyset, \{s_1\}\}$, size 2.
- ▶ $n = 2$: $S = \{s_1, s_2\} \rightsquigarrow \{\emptyset, \{s_1\}, \{s_2\}, \{s_1, s_2\}\}$, size 4.
- ▶ $n = 3$: $S = \{s_1, s_2, s_3\} \rightsquigarrow \left\{ \begin{array}{l} \emptyset, \{s_1\}, \{s_2\}, \{s_1, s_2\}, \\ \{s_3\}, \{s_1, s_3\}, \{s_2, s_3\}, \{s_1, s_2, s_3\} \end{array} \right\}$, 8.

It appears that the number of subsets of S is _____. (notation)

This number also counts _____.

We can label the subsets by whether or not they contain s_i .

For example, for $n = 3$, we label the subsets $\left\{ \begin{array}{l} 000, 100, 010, 110, \\ 001, 101, 011, 111 \end{array} \right\}$

Permutations

Q2. In how many ways can a baseball manager order nine fixed baseball players in a lineup?

Answer: The number of choices for each lineup spot are:

Multiplying gives that the number of lineups is _____ = 362,880.

Definition: A **permutation** of an n -set S is an (ordered) list of **all** elements of S . There are $n!$ such permutations.

Definition: A **k -permutation** of an n -set S is an (ordered) list of k distinct elements of S . **How many are there?**

- ▶ “Permutation” always refers to a list without repetition.

Lists WITHOUT repetition

Question: How many 8-character passwords are there using $A-Z$, $a-z$, $0-9$, containing no repeated character?

OK: 2eas3FGS, 10293465 **Not OK:** 2kdjfn2, oOoOoOo0

Answer: The number of choices for each character are:

for a total of $(62)_8 = \frac{62!}{54!}$ passwords.

In general, the number of words of length k that can be made from an alphabet of length n and where repetition is NOT allowed is $(n)_k$.

- ▶ That is, the number of k -permutations of an n -set is $(n)_k$.
- ▶ **Special case:** For n -permutations of an n -set: $n!$.

Notation

Some quantities appear frequently, so we use shorthand notation:

$$\blacktriangleright [n] := \{1, 2, \dots, n\} \quad \blacktriangleright 2^S := \text{set of all subsets of } S$$

$$\blacktriangleright n! := n \cdot (n-1) \cdot (n-2) \cdots 2 \cdot 1$$

$$\blacktriangleright (n)_k := n \cdot (n-1) \cdot (n-2) \cdots (n-k+1) = \frac{n!}{(n-k)!}$$

$$\blacktriangleright \binom{n}{k} := \frac{n!}{k!(n-k)!} = \frac{(n)_k}{k!}$$

$$\blacktriangleright \binom{\binom{n}{k}}{k} := \binom{k+n-1}{k}$$

★ Leave answers to counting questions in terms of these quantities.

★ **Do NOT** multiply out unless you are comparing values.

Counting subsets of a set

My question: In how many ways are there to choose a subset of k objects out of a set of n objects?

Your answer: $\binom{n}{k}$. “ n choose k ”.

Question: In how many ways can you choose 4 objects out of 10?

Q3. How many Pick-6 lottery tickets are there?
(Choose six numbers between 1–40.)

$$= 3,838,380.$$

- ▶ $\binom{n}{k}$ is called a **binomial coefficient**.
- ▶ Alternate phrasing: How many k -subsets of an n -set are there?
- ▶ The individual objects we are counting are unordered.
They are subsets, not lists.

A formula for $\binom{n}{k}$

You may know that $\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{1}{k!}(n)_k$. But why?

Let's rearrange it.

And prove it!

$$(n)_k = \binom{n}{k} k!$$

We ask the question:

“In how many ways are there to create a k -list of an n -set?”

LHS:

RHS:

Since we counted the same quantity twice, they must be equal!

Counting Multisets

Definition: A **multiset** is an unordered collection of elements where repetition is allowed.

► *Example.* $\{a, a, b, d\}$ is a multiset.

Definition: We say M is a **multisubset** of a set (or multiset) S if every element of M is an element of S .

► *Example.* $M = \{a, a, a, b, d\}$ is a **multisubset** of $S = \{a, b, c, d\}$.

Think Write Pair Share: Enumerate **all** multisubsets of $[3]$.
[In other words, *list them all* or *completely describe the list.*]

Answer:

How would you describe a k -multisubset of $[n]$?


Stars and Bars

Question: How many k -multisets can be made from an n -set?

$$\{a^2, b^0, c^3, d^1\} \quad \begin{array}{l} n = 4 \\ k = 6 \end{array}$$

— *is the same as* —

Question: How many ways are there to place k indistinguishable balls into n distinguishable bins?



— *is the same as* —

Question: How many $\{*, |\}$ -words contain k stars and $(n - 1)$ bars?

** || *** | *

— *which we can count by:* —

Question: How many ways are there to choose k star positions out of $k + n - 1$?

$$\binom{k+n-1}{k} =: \binom{n}{k}$$

Answering Q1–Q4

Q4. How many possible orders for a dozen donuts are there when the store has 30 varieties?

Answer: $\binom{30}{12} = 7,898,654,920$.

Correct order:

| | |
|--|---------------------|
| Q2. Order 9 baseball players ($9!$) | 362,880 |
| Q3. Pick-6; numbers 1–40 $\binom{40}{6}$ | 3,838,380 |
| Q4. 12 donuts from 30 $\binom{30}{12}$ | 7,898,654,920 |
| Q1. 8-character passwords (62^8) | 218,340,105,584,896 |

Summary

| | order matters (choose a list) | order doesn't matter (choose a set) |
|---------------------------|----------------------------------|--|
| repetition allowed | | |
| repetition not allowed | | |