

COMPLEX FENCHEL-NIELSEN COORDINATES WITH SMALL IMAGINARY PARTS

DRAGOMIR ŠARIĆ

ABSTRACT. Kahn and Markovic [9] proved that the fundamental group of each closed hyperbolic three manifold contains a closed surface subgroup. One of the main ingredients in their proof is a theorem which states that an assignment of nearly real, complex Fenchel-Nielsen coordinates to the cuffs of a pants decomposition of a closed surface S induces a quasiFuchsian representation of the fundamental group of S . We give a new proof of this theorem with a slightly stronger conditions on the Fenchel-Nielsen coordinates and explain how to use the exponential mixing of the geodesic flow on a closed hyperbolic three manifold to prove that our theorem is sufficient for the applications in the work of Kahn and Markovic [9].

1. INTRODUCTION

Kahn and Markovic [9] recently proved that the fundamental group $\pi_1(M)$ of any closed hyperbolic three manifold M has a closed surface subgroup. Their proof uses the exponential mixing of the geodesic flow on M in order to find a “well-distributed” finite collection of skew pants in the three manifold M that have large and nearly equal cuff lengths, that are nearly flat, and that can be glued pairwise with nearly zero angles. The collection of skew pants has a subcollection \mathcal{P} that closes to form an abstract closed surface S of (high) genus with nearly real, complex Fenchel-Nielsen coordinates on \mathcal{P} . The final step in the proof of Kahn and Markovic [9] is to show that nearly real, complex Fenchel-Nielsen coordinates on \mathcal{P} necessarily induce an isomorphism between the fundamental group $\pi_1(S)$ and a quasiFuchsian group. Our contribution is to give a new proof of this statement. In fact, we prove a slightly weaker statement by requiring that the imaginary parts of the complex Fenchel-Nielsen coordinates satisfy slightly stronger conditions and establish that this weaker statement is sufficient for the purposes of the proof of the surface subgroup conjecture along the lines in [9]. Our proof adopts the ideas of proving the injectivity of the bending along a measured lamination (cf. [7], [8], [13]) which (at least conceptually) simplifies this part of the argument in [9].

Let S be a closed surface of genus $g \geq 2$ equipped with a pants decomposition \mathcal{P} . Then \mathcal{P} consists of $3g - 3$ simple closed curves such that each component of the complement is a pair of pants. Following [9, §2] (see also §2), to each cuff $C \in \mathcal{P}$ we associate complex half-length $hl(C) \in \mathbb{C}/2\pi i\mathbb{Z}$ and complex twist-bend parameter $s(C) \in \mathbb{C}/(2\pi i\mathbb{Z} + hl(C)\mathbb{Z})$. An assignment of half-lengths and twist-bend parameters to $C \in \mathcal{P}$ induces a representation of the fundamental group $\pi_1(S)$ into $PSL_2(\mathbb{C})$. The representation is Fuchsian if and only if $\{(hl(C), s(C))\}_{C \in \mathcal{P}} \in \mathbb{R}^{3g-3}$.

Date: April 27, 2012.

This research is partially supported by National Science Foundation grant DMS 1102440.

The following theorem characterizes a neighborhood of the real subspace \mathbb{R}^{3g-3} inside \mathbb{C}^{3g-3} which gives quasiFuchsian representations. We note that the size of the neighborhood is independent of the genus g .

Theorem 1.1 (Kahn-Markovic [9]). *There exist universal $\hat{\epsilon}, K_0 > 0$ and $R(\hat{\epsilon}) > 0$ such that the following is satisfied. Let S be a closed surface of genus $g \geq 2$ and let \mathcal{P} be a pants decomposition of S . If $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$ is a representation which is discrete and faithful on each pair of pants in \mathcal{P} and if the reduced complex Fenchel-Nielsen coordinates on each cuff $C \in \mathcal{P}$ satisfy*

$$(1) \quad |hl(C) - R/2| < \epsilon$$

and

$$(2) \quad |s(C) - 1| < \epsilon/R$$

for some $\epsilon < \hat{\epsilon}$ and $R > R(\hat{\epsilon})$, then $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$ is injective and $\rho(\pi_1(S))$ is quasiFuchsian.

Moreover, let S be endowed with a hyperbolic metric whose reduced Fenchel-Nielsen coordinates are $hl(C) = R/2$ and $s(C) = 1$ for each $C \in \mathcal{P}$. Then there exists an injective map $\tilde{f} : \partial_\infty \tilde{S} \rightarrow \partial_\infty \mathbb{H}^3$ which conjugates $\pi_1(S)$ into the above quasiFuchsian group and which extends to a $(1 + K_0\epsilon)$ -quasiconformal map of $\partial_\infty \mathbb{H}^3$ onto itself.

We give a new proof of the above theorem under assumptions (2) and

$$(3) \quad |hl(C) - R/2| < \epsilon/R.$$

Even though (3) is stronger than (1) (which makes our statement weaker than the above theorem), it turns out that this is enough for the purposes in [9]. At the end of Introduction we indicate how to see that (3) follows from the fact that the skew pairs of pants are “well-distributed” inside the three manifold which proves that the weaker statement suffices. One advantage of using (3) instead of (1) is that we do not need to require that ρ is discrete and faithful on pairs of pants in \mathcal{P} in order to establish the injectivity of the representation $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$. In particular, ρ is discrete and injective on each pair of pants of \mathcal{P} if it satisfies (2) and (3).

In [9], Theorem 1.1 is proved by estimating the derivative (along a path of representations connecting the Fuchsian representation with the representation corresponding to the given reduced Fenchel-Nielsen coordinates) of the distance between the images in \mathbb{H}^3 of two lifts of geodesics $C \in \mathcal{P}$ in \mathbb{H}^2 from the above by a function of the distance between these two lifts of geodesics at the representation. This leads to an inductive argument which gives the desired theorem.

Our approach is to decompose each pair of pants in \mathcal{P} into two ideal hyperbolic triangles by adding three infinite geodesics such that each end of each added geodesic spirals around a different cuff. The union of cuffs of \mathcal{P} together with the added geodesics in each pair of pants is a maximal geodesic lamination λ in S with finitely many leaves. Let $\tilde{\lambda}$ be the lift of λ to the universal covering \mathbb{H}^2 . The reduced complex Fenchel-Nielsen coordinates $\{(hl(C), s(C))\}_{C \in \mathcal{P}}$ induce a developing map $\tilde{f} : \partial_\infty \mathbb{H}^2 \rightarrow \partial_\infty \mathbb{H}^3$ which conjugates $\pi_1(S) < PSL_2(\mathbb{C})$ into a subgroup of $PSL_2(\mathbb{C})$. The developing map extends to complementary triangles of $\tilde{\lambda}$ to define a pleated surface $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ which is pleated along $\tilde{\lambda}$ (cf. [3] and also §3). Each

pleated surface along $\tilde{\lambda}$ induces a finitely additive $(\mathbb{C}/2\pi i\mathbb{Z})$ -valued transverse cocycle α to $\tilde{\lambda}$ which measures the shearing and the bending along $\tilde{\lambda}$ (cf. [3]). The *bending cocycle* is the imaginary part β of the transverse cocycle α which is an $(\mathbb{R}/2\pi\mathbb{Z})$ -valued transverse cocycle to $\tilde{\lambda}$ measuring the amount of the bending of the pleated surface. We translate the reduced complex Fenchel-Nielsen coordinates into the bending transverse cocycle β to $\tilde{\lambda}$ as follows.

An isolated leaf \tilde{l} of $\tilde{\lambda}$ is on a common boundary of two complementary ideal triangles $\Delta^1(\tilde{l})$ and $\Delta^2(\tilde{l})$ to $\tilde{\lambda}$. Isolated leaves of $\tilde{\lambda}$ accumulate to each lift \tilde{C} of each cuff $C \in \mathcal{P}$ from both sides of \tilde{C} . Let S be endowed with a hyperbolic metric whose Fenchel-Nielsen coordinates are $\{(Re(hl(C)), Re(s(C)))\}_{C \in \mathcal{P}}$ and divide each pair of pants of \mathcal{P} into two hyperbolic hexagons Σ_1 and Σ_2 by drawing common orthogonal arcs between pairs of cuffs of each pair of pants in \mathcal{P} . Each hexagon Σ on the surface S lifts to infinitely many hexagons in the universal covering \mathbb{H}^2 . Recall the assumptions $|Re(s(C)) - 1| < \epsilon/R$ and $|Re(hl(C)) - R/2| < \epsilon/R$, for $R \geq R(\hat{\epsilon})$ and $0 < \epsilon < \hat{\epsilon}$. A cuff C is the union of two boundary sides of two hexagons coming from the pair of pants on one side of C as well as the union of two boundary sides of two hexagons coming from the pair of pants on the other side of C . The boundary sides of the hexagons from one side of C are not exactly matched along C with the boundaries of the hexagons from the other side of C but they are glued with a shift close to 1 by the condition $|Re(s(C)) - 1| < \epsilon/R$. It follows that for each hexagon Σ_i , $i = 1, 2$, on one side of C there is a unique hexagon Σ'_i , $i = 1, 2$, on the other side of C such that the common subarc of their boundary sides on C has length close to $R/2 - 1$. We will say that Σ_i and Σ'_i are 0-neighbors in this case. Two lifts $\tilde{\Sigma}_i$ and $\tilde{\Sigma}'_i$ to \mathbb{H}^2 of 0-neighbors hexagons are also called 0-neighbors if they meet along a lift \tilde{C} of C with a common subarc of length close to $R/2 - 1$ (cf. §3). If hexagon $\tilde{\Sigma}$ is a lift of a hexagon Σ , then $\tilde{\Sigma}$ intersects infinitely many complementary triangles to $\tilde{\lambda}$. There is a unique triangle $\Delta_{\tilde{\Sigma}}$ such that its intersection with $\tilde{\Sigma}$ is a hexagon, and we call $\Delta_{\tilde{\Sigma}}$ the *canonical triangle* of $\tilde{\Sigma}$ (cf. §3 and Figure 1).

Theorem 1.2. *There exists $C_0 > 0$ such that the following holds. Let $\{hl(C), s(C)\}_{C \in \mathcal{P}}$ be the reduced complex Fenchel-Nielsen coordinates that satisfy (3) and (2), and let β be the induced bending transverse cocycle to the lamination $\tilde{\lambda}$. If \tilde{l} is an isolated leaf of $\tilde{\lambda}$ and $\Delta^i(\tilde{l})$, $i = 1, 2$, are complementary triangles to $\tilde{\lambda}$ with a common boundary side \tilde{l} , then*

$$(4) \quad |\beta(\Delta^1(\tilde{l}), \Delta^2(\tilde{l}))| \leq \frac{C_0 \epsilon}{R}.$$

If $\tilde{\Sigma}_1$ and $\tilde{\Sigma}_2$ are 0-neighbors hexagons, and $\Delta_{\tilde{\Sigma}_1}$ and $\Delta_{\tilde{\Sigma}_2}$ are their canonical triangles, then

$$(5) \quad |\beta(\Delta_{\tilde{\Sigma}_1}, \Delta_{\tilde{\Sigma}_2})| \leq \frac{C_0 \epsilon}{R}.$$

Theorem 1.2 translates the original problem of whether the reduced complex Fenchel-Nielsen coordinates give a quasiFuchsian representation into whether a bending cocycle gives a quasiFuchsian representation. We point out that any condition on the bending cocycle that guarantees injectivity of the bending map on $\partial_\infty \mathbb{H}^2$ necessarily depends on the hyperbolic metric from which the bending starts. In our case, a sufficient information about the hyperbolic metric is given by the fact

that a unit length hyperbolic arc in \mathbb{H}^2 can have at most $2R + 2$ intersections with the lifts of the cuffs (cf. [9, Lemma 2.3] and Lemma 3.2). When the transverse bending measure is countably additive, then it is well-known that the bending measure which is uniformly small on each transverse arc of length 1 gives a bending map which is injective on $\partial_\infty\mathbb{H}^2$ (cf. [7], [8], [13]). The above sufficient condition for the injectivity of the bending along the measured laminations is universal in the sense that it does not depend on the genus of the surface and, in fact, it works for any surface including the unit disk. The main difficulty in proving that the bending along finitely additive transverse cocycles is injective on $\partial_\infty\mathbb{H}^2$ lies in the fact that the total variation of the transverse bending measure is infinite. However, conditions (3) and (2) guarantee that the bending map behaves “semi-locally” as the bending map along a measured lamination. Using ideas from [13], we prove that the above conditions on the bending cocycle and the hyperbolic metric are sufficient to guarantee injectivity on $\partial_\infty\mathbb{H}^2$ of the bending map. Holomorphic motions provide the desired bound on quasiconformal extension of the restriction to $\partial_\infty\mathbb{H}^2$ of the bending map.

Theorem 1.3. *Given C_0 , there exist $\hat{\epsilon} > 0$, $K_0 > 1$ and $R(\hat{\epsilon}) > 0$ such that for $0 \leq \epsilon < \hat{\epsilon}$ the following is satisfied. Let S be a closed hyperbolic surface equipped with a maximal, finite geodesic lamination λ such that each closed geodesic of λ has length in the interval $(R - \frac{C_0\epsilon}{R}, R + \frac{C_0\epsilon}{R})$ for some $R \geq R(\hat{\epsilon})$ and that each geodesic arc on S of length 1 intersects at most $C_0 \cdot R$ closed geodesics of λ . If a bending cocycle β transverse to the lift $\tilde{\lambda}$ in \mathbb{H}^2 satisfies (4) and (5) then the induced bending map*

$$\tilde{f}_\beta : \partial_\infty\mathbb{H}^2 \rightarrow \partial_\infty\mathbb{H}^3$$

is injective and the induced representation of $\pi_1(S)$ is quasiFuchsian. The bending map extends to a $(1 + K_0\epsilon)$ -quasiconformal map $f_\beta : \partial_\infty\mathbb{H}^3 \rightarrow \partial_\infty\mathbb{H}^3$.

We give an analogue of the above theorem for non-finite geodesic laminations and bending (finitely additive) transverse cocycles [14].

It remains to explain why the condition $|hl(C) - R/2| < \epsilon$ can be replaced with the condition $|hl(C) - R/2| < \epsilon/R$. The geodesic flow on a closed hyperbolic three manifold M is exponentially mixing [12]. Let $\mathcal{F}(M)$ be the 2-frame bundle over M . Since the hyperbolic Laplacian of M has a spectral gap the following holds by [12]. There exists $q > 0$ which depends on the three manifold M such that for any two C^{mifty} -functions $\psi, \varphi : \mathcal{F}(M) \rightarrow \mathbb{R}$ we have

$$(6) \quad \left| \Lambda(\mathcal{F}(M)) \int_{\mathcal{F}(M)} (g_t^* \psi) \varphi d\Lambda - \int_{\mathcal{F}(M)} \psi d\Lambda \int_{\mathcal{F}(M)} \varphi d\Lambda \right| \leq C e^{-qt},$$

where Λ is the Liouville measure on \mathcal{F} and the constant C depends on the C^1 -norms of ψ and φ .

Let $f_\epsilon : \mathcal{F}(M) \rightarrow \mathbb{R}$ be a non-negative C^∞ -function supported in the ϵ -neighborhood of a point in $\mathcal{F}(M^3)$ with $\int_{\mathcal{F}(M)} f_\epsilon d\Lambda = 1$, called a *bump function* for the ϵ -neighborhood. By applying (6) to f_ϵ , Kahn and Markovic [9] proved that there exist triples of 2-frames, called *tripods*, that after traveling a long time $t > 0$ along the geodesic flow return to their ϵ -neighborhoods. These tripods define skew pairs of pants in M whose cuff lengths are $R = 2t - 2 \log \frac{4}{3}$ with a possible error $D \cdot \epsilon$, for some $D > 0$ because the expression on the right of (6) goes to 0 as $t \rightarrow \infty$ (cf. [9, Lemma 4.6]).

It is possible to improve the estimate on the complex length of the cuffs of the above skew pairs of pants. Note that the constant C in (6) can be estimated (cf. [12]) in terms of H_2^2 -Sobolev norms to be $C_0 \|\psi\|_{H_2^2} \|\varphi\|_{H_2^2}$, where C_0 is a fixed constant. Then, for a given time t geodesic flow, we consider bump function $f_{\epsilon/t}$ for ϵ/t -neighborhood of a point in $\mathcal{F}(M)$. The bump function $f_{\epsilon/t}$ can be produced by scaling the domain and the size of f_ϵ such that $\|f_{\epsilon/t}\|_{H_2^2} \leq p_2(t) \|f_\epsilon\|_{H_2^2}$, where $p_2(t)$ is a polynomial in t of degree 2. From (6) we get

$$\left| \Lambda(\mathcal{F}(M)) \int_{\mathcal{F}(M)} (g_t^* f_{\epsilon/t}) f_{\epsilon/t} d\Lambda - 1 \right| \leq C_0 \|f_\epsilon\|_{H_2^2}^2 p_2(t)^2 e^{-qt} \rightarrow 0$$

as $t \rightarrow \infty$. This implies that the skew pairs of pants have cuffs of the length R within ϵ/R , when we choose an appropriate value for $t = t(R)$ thus obtaining (3).

Acknowledgements. I am grateful to Vladimir Markovic and Hideki Miyachi for various discussions regarding this work.

2. THE REDUCED COMPLEX FENCHEL-NIELSEN COORDINATES

Let S be a closed surface of genus $g \geq 2$ and let $\pi_1(S)$ be its fundamental group. Let \mathcal{P} be a pants decomposition of S , namely \mathcal{P} consists of $3g - 3$ simple, closed curves on S such that the components of the complement of \mathcal{P} are pairs of pants. A representation $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$ associates to each cuff $C \in \mathcal{P}$ two complex numbers: the complex length and the twist-bend parameter. In total, $6g - 6$ complex numbers is associated to a representation ρ , called the complex Fenchel-Nielsen coordinates. The complex Fenchel-Nielsen coordinates were introduced in [10] and [16], and it was proved there that the quasiFuchsian space of S is parametrized by an open subset of \mathbb{C}^{6g-6} which contains the real locus \mathbb{R}^{6g-6} . We use the reduced Fenchel-Nielsen coordinates introduced by Kahn and Markovic [9, §2] and we refer the reader to their article for more details.

Let α and β be two oriented geodesics in \mathbb{H}^3 . Let γ be their common orthogonal oriented from α to β . The complex distance $d_\gamma(\alpha, \beta)$ between α and β is defined to have a positive real part equal to the distance between $\alpha \cap \gamma$ and $\beta \cap \gamma$, while the imaginary part of $d_\gamma(\alpha, \beta)$ is the angle between the parallel transport along γ of the unit tangent vector to α at $\alpha \cap \gamma$ and the unit tangent vector to β at $\beta \cap \gamma$. Since the imaginary part of $d_\gamma(\alpha, \beta)$ is well defined modulo $2\pi i$, we have $d_\gamma(\alpha, \beta) \in \mathbb{C}/2\pi i\mathbb{Z}$ (for more details, cf. [9, §2]).

Let Π_0 be a pair of pants and $\pi_1(\Pi_0)$ be its fundamental group. Consider a representation $\rho : \pi_1(\Pi_0) \rightarrow PSL_2(\mathbb{C})$ which is faithful and loxodromic. Namely, the cuffs C_i , $i = 0, 1, 2$, of Π_0 are represented by loxodromic elements $\rho(C_i) \in PSL_2(\mathbb{C})$. Let γ_i be the axis of $\rho(C_i)$ and η_i be the common orthogonal to γ_{i-1} and γ_{i+1} , for $i = 0, 1, 2$, where the indices are taken modulo 3. Then the half-length $hl_{\Pi_0, \rho}(C_i)$ of the curve C_i associated to the representation ρ is $d_{\gamma_i}(\eta_{i-1}, \eta_{i+1})$ (cf. [9, §2]).

Consider a representation $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$ of the fundamental group $\pi_1(S)$ of a closed surface S of genus at least two into $PSL_2(\mathbb{C})$ and fix a pants decomposition \mathcal{P} . The representation ρ is *viable* if $\rho : \pi_1(\Pi) \rightarrow PSL_2(\mathbb{C})$ is discrete and faithful for each pair of pants Π of the pants decomposition \mathcal{P} , and for any two pairs of pants Π and Π' with a common cuff C we have $hl_{\Pi, \rho}(C) = hl_{\Pi', \rho}(C)$ (cf. [9]). For a given viable representation ρ , we define the complex half-length of a cuff $C \in \mathcal{P}$ by $hl(C) = hl_{\Pi, \rho}(C) = hl_{\Pi', \rho}(C) \in \mathbb{C}/2\pi i\mathbb{Z}$, where Π and Π' are pairs

of pants with one cuff C (cf. [9]). Let Π and Π' be two pairs of pants with cuffs C_i , $i = 0, 1, 2$, and C'_i , $i = 0, 1, 2$, respectively such that $C_0 = C'_0 = C$. Let γ_i, γ'_i be the axes of $\rho(C_i), \rho(C'_i)$. Let η_i and η'_i be common orthogonals to $\gamma_{i-1}, \gamma_{i+1}$ and $\gamma'_{i-1}, \gamma'_{i+1}$, respectively. The *twist-bend parameter* $s(C)$ is the complex distance between unit tangent vector to η_1 at the point $\eta_1 \cap \gamma_0$ and the unit tangent vector to η'_1 at the point $\eta'_1 \cap \gamma_0$. The choices involved guarantee that the complex twist-bend parameter $s(C)$ is well defined in $\mathbb{C}/(2\pi i\mathbb{Z} + hl(C)\mathbb{Z})$.

3. PLEATED SURFACES AND TRANSVERSE COCYCLES TO GEODESIC LAMINATIONS

Recall that S is a closed surface of genus $g \geq 2$. Let λ be a *maximal* geodesic lamination on S , namely each component of the complement of λ is an ideal hyperbolic triangle. We do not need to specify a hyperbolic metric on S in order to be able to talk about geodesic laminations on S (cf. [3]).

Let $\pi : \mathbb{H}^2 \rightarrow S$ be the universal covering for a metric m and let $\tilde{\lambda} = \pi^{-1}(\lambda)$. An *abstract pleated surface* for S with the pleating locus λ is a pleating map \tilde{f} with the pleating locus $\tilde{\lambda}$ from the hyperbolic plane \mathbb{H}^2 into the hyperbolic three-space \mathbb{H}^3 which is equivariant under the action on \mathbb{H}^2 of the covering group G of S and the action on \mathbb{H}^3 of a subgroup $G_{\tilde{f}}$ of $PSL_2(\mathbb{C})$. If the continuous extension of \tilde{f} from the ideal boundary $\partial_\infty \mathbb{H}^2$ of \mathbb{H}^2 to the ideal boundary $\partial_\infty \mathbb{H}^3$ is injective then the group $G_{\tilde{f}}$ is quasifuchsian and \tilde{f} projects to a pleated map from $S = \mathbb{H}^2/G$ into the quasifuchsian three-manifold $\mathbb{H}^3/G_{\tilde{f}}$.

In this article we consider only finite maximal geodesic laminations on S which are necessarily obtained by triangulating pairs of pants of a pants decomposition \mathcal{P} of S as follows. Let Π_1 and Π_2 be two pairs of pants of \mathcal{P} that have $C \in \mathcal{P}$ as one of its boundaries. It is possible that $\Pi_1 = \Pi_2$. Assume that ideal triangulations of Π_1 and Π_2 are given. Let a_1^j for $j = 1, 2$ be the boundary edges of the triangulation of Π_1 whose one end accumulate at C , and similarly let a_2^j for $j = 1, 2$ be the boundary edges of the triangulation of Π_2 whose one end accumulate at C . Let \tilde{C} be a lift of C to \mathbb{H}^2 . Then there are adjacent lifts \tilde{a}_1^j of a_1^j for $j = 1, 2$ that share an ideal endpoint x_1 with \tilde{C} , and there are adjacent lifts \tilde{a}_2^j of a_2^j for $j = 1, 2$ that share an ideal endpoint x_2 with \tilde{C} . Either $x_1 = x_2$ or $x_1 \neq x_2$. If $x_1 = x_2$ then, we say that the triangulations of the two pairs of pants with common boundary C *accumulate in the same direction on C* . From now on, we assume that $\lambda_{\mathcal{P}}$ is a finite, maximal geodesic lamination that is obtained by triangulating pairs of pants of \mathcal{P} such that the triangulations of pairs of pants with common boundaries accumulate in the same direction at each $C \in \mathcal{P}$.

Let $\{\Pi_j\}_{j=1}^{2g-2}$ be the pairs of pants in \mathcal{P} . Given a pair of pants Π_j in \mathcal{P} with cuffs $C_i^j \in \mathcal{P}$, $i = 1, 2, 3$, denote by $\gamma_i^j \in \pi_1(S)$ the elements representing closed curves C_i^j such that $\gamma_3^j \gamma_2^j \gamma_1^j = id$ in $\pi_1(S)$. Let $\{(hl(C), s(C))\}_{C \in \mathcal{P}}$ be the reduced complex Fenchel-Nielsen coordinates that satisfy (2) and (3). By [10, Proposition 2.3], there exists a representation $\rho : \pi_1(S) \rightarrow PSL_2(\mathbb{C})$ which realizes $\{(hl(C), s(C))\}_{C \in \mathcal{P}}$ such that $\rho(\gamma_i^j) \in PSL_2(\mathbb{C})$, for $i = 1, 2, 3$, and $j = 1, 2, \dots, 2g-2$ are loxodromic and $\rho(\gamma_i^j)$, $i = 1, 2, 3$, have distinct endpoints for each $j = 1, 2, \dots, 2g-2$. Let S be endowed with the hyperbolic metric whose Fenchel-Nielsen coordinates $\{(Re(hl(C)), Re(s(C)))\}_{C \in \mathcal{P}}$ are the real parts of the reduced complex Fenchel-Nielsen coordinates $\{(hl(C), s(C))\}_{C \in \mathcal{P}}$. Let G be the covering group for the universal covering $\pi : \mathbb{H}^2 \rightarrow S$. Consider the lifts \tilde{C} to \mathbb{H}^2 of the cuffs of \mathcal{P} . Then there

exists a developing map \tilde{f} from the set of endpoints of $\tilde{C} \in \mathcal{P}$ into $\partial_\infty \mathbb{H}^3$ which realizes the reduced complex Fenchel-Nielsen coordinates $\{hl(C), s(C)\}_{C \in \mathcal{P}}$. The map \tilde{f} extends to an abstract pleating map as follows. Let $\tilde{\lambda}_{\mathcal{P}}$ be the lift of $\lambda_{\mathcal{P}}$ to \mathbb{H}^2 . Each geodesic of $\tilde{\lambda}_{\mathcal{P}}$ which is not a lift of a cuff has its both endpoints at the endpoints of two lifts of two different cuffs of a single pair of pants in \mathcal{P} which implies that the endpoints are distinct. Thus \tilde{f} extends to map each geodesic of $\tilde{\lambda}_{\mathcal{P}}$ into a geodesic of \mathbb{H}^3 . Since the complementary components to $\tilde{\lambda}_{\mathcal{P}}$ are ideal hyperbolic triangles, it follows that we have an extension $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ which determines an abstract pleated surface with the pleating locus $\lambda_{\mathcal{P}}$. Thus the representation ρ realizes the geodesic lamination $\tilde{\lambda}_{\mathcal{P}}$ (cf. [2]).

An abstract pleated surface $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ with a pleating locus $\tilde{\lambda}_{\mathcal{P}}$ determines a $(\mathbb{C}/2\pi i\mathbb{Z})$ -valued transverse cocycle α to the geodesic lamination $\tilde{\lambda}_{\mathcal{P}}$ (cf. [3]). Namely, α determines a finitely additive assignment of a number in $\mathbb{C}/2\pi i\mathbb{Z}$ to each arc transverse to $\tilde{\lambda}_{\mathcal{P}}$ (with endpoints in the complementary triangles of $\tilde{\lambda}_{\mathcal{P}}$) which is homotopy invariant relative $\lambda_{\mathcal{P}}$. If k is a geodesic arc connecting triangles Δ_1 and Δ_2 , then we write $\alpha(\Delta_1, \Delta_2) = \alpha(k)$ because $\alpha(k)$ depends only on the homotopy class of k relative $\tilde{\lambda}_{\mathcal{P}}$. The real part of α is an \mathbb{R} -valued transverse cocycle which completely determines the path metric on the pleated surface (cf. [3]). The imaginary part of α is an $(\mathbb{R}/2\pi\mathbb{Z})$ -valued transverse cocycle β to the geodesic lamination $\lambda_{\mathcal{P}}$. The transverse cocycle β determines the amount of the bending of the pleated surface $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ (cf. [3]).

Our first task is to translate the conditions (3) and (2) in terms of the associated transverse cocycle to $\tilde{\lambda}_{\mathcal{P}}$. Let $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ be the pleating map corresponding to the reduced Fenchel-Nielsen coordinates $\{hl(C), s(C)\}_{C \in \mathcal{P}}$ starting from the real Fenchel-Nielsen coordinates $\{Re(hl(C)), Re(s(C))\}_{C \in \mathcal{P}}$ as above. We also lift the decomposition of the pairs of pants of \mathcal{P} into right-angled hexagons. A right-angled hexagons on S lifts to an infinite collection of right-angled hexagons in \mathbb{H}^2 . Fix a lift $\tilde{C} \in \pi^{-1}(C)$ of the closed geodesic C and fix a lifted hexagon Σ that has one boundary side on \tilde{C} . Then there is a unique lifted hexagon Σ' with one boundary side on \tilde{C} which lies on the opposite side of \tilde{C} such that the distance between the corresponding vertices of Σ and Σ' on \tilde{C} is equal to $Re(s(C))$. We say that Σ and Σ' are *0-neighbors*. Thus 0-neighbors hexagons meet along \tilde{C} and have an arc of length $R/2 - Re(s(C))$ in common (cf. Figure 1). If two hexagons Σ and Σ'' meet along their boundaries but they are not 0-neighbors then we call them *1-neighbors* (cf. Figure 1).

Fix a lifted hexagon Σ in \mathbb{H}^2 . Among all complementary triangles to $\tilde{\lambda}_{\mathcal{P}}$ there is a unique triangle Δ_Σ whose all three boundary sides intersect Σ . We call Δ_Σ the *canonical triangle* for Σ . Let Σ_t be the intersection of Σ and Δ_Σ . Then $\Sigma \setminus \Sigma_t$ has three connected components each being a quadrilateral (cf. Figure 1). Let \mathcal{H} be the set of all lifted hexagons in \mathbb{H}^2 . Then

$$\mathcal{TH}_t = \bigcup_{\Sigma \in \mathcal{H}} \Sigma_t$$

separates geodesics in $\pi^{-1}(C)$ (cf. Figure 2). We give a proof of Theorem 1.2 from Introduction.




FIGURE 1. Σ and Σ' are 0-neighbors; Σ and Σ'' are 1-neighbors; Δ_Σ is the characteristic triangle of Σ




FIGURE 2. The set \mathcal{TH}_t

Theorem 3.1. *There exists $C_0 > 0$ such that the following holds. Let $\{hl(C), s(C)\}_{C \in \mathcal{P}}$ be the reduced complex Fenchel-Nielsen coordinates such that*

$$|hl(C) - R/2| < \frac{\epsilon}{R}$$

and

$$|s(C) - 1| < \frac{\epsilon}{R}.$$

Let S be endowed with a hyperbolic metric whose Fenchel-Nielsen coordinates are $\{\operatorname{Re}(hl(C)), \operatorname{Re}(s(C))\}_{C \in \mathcal{P}}$. Let $\lambda_{\mathcal{P}}$ be a maximal geodesic lamination obtained by triangulating pairs of pants of \mathcal{P} such that the edges of the triangles from both sides of each $C \in \mathcal{P}$ accumulate in the same direction. Let $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^3$ be the bending

map with the bending locus $\tilde{\lambda}_{\mathcal{P}} = \pi^{-1}(\lambda_{\mathcal{P}})$ which realizes the complex Fenchel-Nielsen coordinates $\{hl(C), s(C)\}_{C \in \mathcal{P}}$, where $\pi : \mathbb{H}^2 \rightarrow S$ is the universal covering. Denote by β the bending cocycle transverse to $\tilde{\lambda}_{\mathcal{P}}$ for \tilde{f} . Let \tilde{l} be an isolated leaf of $\tilde{\lambda}_{\mathcal{P}}$ which is on the boundary of two complementary triangles $\Delta_1(\tilde{l})$ and $\Delta_2(\tilde{l})$ of $\tilde{\lambda}_{\mathcal{P}}$. Then

$$|\beta(\Delta_1(\tilde{l}), \Delta_2(\tilde{l}))| \leq \frac{C_0 \epsilon}{R}.$$

Moreover, let Σ_1 and Σ_2 be 0-neighbor hexagons and let Δ_{Σ_1} and Δ_{Σ_2} be their canonical triangles, respectively. Then

$$|\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma_2})| \leq \frac{C_0 \epsilon}{R}.$$

Proof. Let Π be a pair of pants in \mathcal{P} with boundary curves C_i , $i = 0, 1, 2$. Let l_i , for $i = 0, 1, 2$, be the geodesics of $\lambda_{\mathcal{P}}$ that triangulate Π such that l_{i+1} and l_{i+2} accumulate on C_i , for $i = 0, 1, 2$, with the indices taken modulo 3. Let \tilde{C}_i be a lift of C_i to \mathbb{H}^2 and let $\tilde{l}_{i+1}, \tilde{l}_{i+2}$ be consecutive lifts of l_{i+1}, l_{i+2} that share a common endpoint with \tilde{C}_i . Let $\gamma_i \in PSL_2(\mathbb{R})$ be the deck transformation corresponding to \tilde{C}_i such that $\gamma_i(\tilde{l}_{i+1}) = \tilde{l}'_{i+1}$ is adjacent to \tilde{l}_{i+2} . Let $\tilde{l}'_{i+2} = \gamma_i(\tilde{l}_{i+2})$. Then \tilde{l}'_{i+2} is adjacent to \tilde{l}'_{i+1} .

Let r_{i+2} be the geodesic ray orthogonal to $\tilde{f}(\tilde{l}_{i+2})$ that starts at the endpoint of $\tilde{f}(\tilde{l}_{i+1})$ which is not in common with $\tilde{f}(\tilde{C}_i)$. Let r'_{i+2} be the geodesic ray orthogonal to $\tilde{f}(\tilde{l}_{i+2})$ that starts at the endpoint of $\tilde{f}(\tilde{l}'_{i+1})$ which is not in common with $\tilde{f}(\tilde{C}_i)$. Define s_{i+2} to be the complex distance between the unit tangent vector to r'_{i+2} at $r'_{i+2} \cap \tilde{f}(\tilde{l}_{i+2})$ and the unit tangent vector to r_{i+2} at $r_{i+2} \cap \tilde{f}(\tilde{l}_{i+2})$. Define s_{i+1} using the geodesics $\tilde{f}(\tilde{l}_{i+2}), \tilde{f}(\tilde{l}'_{i+1}), \tilde{f}(\tilde{l}_{i+2})$ similar to the above. Then

$$s_{i+1} + s_{i+2} = l(\delta_i)$$

where $\delta_i = \tilde{f} \circ \gamma_i \circ \tilde{f}^{-1} \in PSL_2(\mathbb{C})$ and $l(\delta_i)$ is the complex translation length of δ_i , for $i = 0, 1, 2$. Solving the above system gives

$$s_i = \frac{l(\delta_{i+1}) + l(\delta_{i+2}) - l(\delta_i)}{2}$$

for $i = 0, 1, 2$. Since

$$|Im(\frac{1}{2}l(\delta_i))| \leq \frac{\epsilon}{R},$$

it follows that

$$|\beta(\Delta_1(\tilde{l}_i), \Delta_2(\tilde{l}_i))| \leq \frac{C_0 \epsilon}{R}$$

for $i = 0, 1, 2$ and some $C_0 > 0$.

Let $\Sigma \subset \mathbb{H}^2$ be a lifted right angled hexagon from a pair of pants Π whose boundary sides lie on C_i , $i = 0, 1, 2$. We fix lifts \tilde{C}_i of C_i such that three sides of Σ lie on \tilde{C}_i for $i = 0, 1, 2$. Let $\Sigma_{\tilde{f}}$ be the skew right angled hexagon whose three sides lie on $\tilde{f}(C_1), \tilde{f}(C_2)$ and $\tilde{f}(C_3)$. Then the complex lengths of these sides are $hl(C_1), hl(C_2)$ and $hl(C_3)$. These sides are called *long sides* and the other three sides of $\Sigma_{\tilde{f}}$ are called *short sides*. Denote by h_i the short side of $\Sigma_{\tilde{f}}$ which connects $\tilde{f}(C_{i+1})$ and $\tilde{f}(C_{i+2})$. Then the hexagon cosine formula directly gives (cf. [4], [9])

$$l(h_i) = 2e^{-\frac{R}{4} + \frac{1}{2}[hl(C_{i+1}) + hl(C_{i+2}) - hl(C_i)]} + O(e^{-3R/4})$$




FIGURE 3. Computing the bending cocycle.

where $l(h_i)$ is the complex distance between $\tilde{f}(\tilde{C}_{i+1})$ and $\tilde{f}(\tilde{C}_{i+2})$. This implies

$$|\operatorname{Re}(l(h_i))| = O(e^{-R/4})$$

and

$$|\operatorname{Im}(l(h_i))| = O\left(\frac{\epsilon}{R}e^{-R/4}\right).$$

Let Σ_1 and Σ_2 be two 0-neighbors hexagons in \mathbb{H}^2 , and let Δ_{Σ_1} and Δ_{Σ_2} be their canonical triangles. Let $\tilde{C} \in \tilde{\mathcal{P}} = \pi^{-1}(\mathcal{P})$ be the geodesic which separates Δ_{Σ_1} and Δ_{Σ_2} , and let $C = \pi(\tilde{C}) \in \mathcal{P}$. Note that both Σ_1 and Σ_2 have one boundary side on \tilde{C} . Normalize the bending map such that the ideal triangles $\tilde{f}(\Delta_{\Sigma_1})$ and $\tilde{f}(\Delta_{\Sigma_2})$, have a common endpoint ∞ , and that $\tilde{f}(\tilde{C})$ has endpoints 0 and ∞ . Let \tilde{C}_1^j for $j = 1, 2$ be the two geodesics of $\tilde{\mathcal{P}}$ (different from \tilde{C}) which contain boundary sides of Σ_1 , and let \tilde{C}_2^j for $j = 1, 2$ be the two geodesics in $\tilde{\mathcal{P}}$ (different from \tilde{C}) which contain boundary sides of Σ_2 . We can assume that the twist-bend $s(C)$ is the complex distance (along $\tilde{f}(\tilde{C})$) between the common orthogonal to $\tilde{f}(\tilde{C}_1^1)$ and $\tilde{f}(\tilde{C})$, and the common orthogonal to $\tilde{f}(\tilde{C}_2^1)$ and $\tilde{f}(\tilde{C})$. It follows that the complex distance (along $\tilde{f}(\tilde{C})$) between the common orthogonal to $\tilde{f}(\tilde{C}_1^2)$ and $\tilde{f}(\tilde{C})$, and the common orthogonal to $\tilde{f}(\tilde{C}_2^2)$ and $\tilde{f}(\tilde{C})$ is also equal to the twist-bend $s(C)$ (cf. Figure 3).

We recall the definition of $\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma_2})$ given by Bonahon [3]. Let \mathcal{W} be the component of $\mathbb{H}^2 \setminus (\Delta_{\Sigma_1} \cup \Delta_{\Sigma_2})$ which separates Δ_{Σ_1} and Δ_{Σ_2} . Denote by $\tilde{\lambda}_{\mathcal{P}}(\Delta_{\Sigma_1}, \Delta_{\Sigma_2})$ the set of leaves of $\tilde{\lambda}_{\mathcal{P}}$ which separate Δ_{Σ_1} and Δ_{Σ_2} , and orient them to the left as seen from Δ_{Σ_1} . The leaves of $\tilde{\lambda}_{\mathcal{P}}$ divide \mathcal{W} into hyperbolic strips and the images under \tilde{f} of the hyperbolic strips are two-dimensional hyperbolic strips in \mathbb{H}^3 . Each such hyperbolic strip intersects $\partial_{\infty}\mathbb{H}^3$ in two circular arcs with a possibility that one is reduced to a point such that one circular arc is bounded by the negative endpoints of the leaves $\tilde{f}(\tilde{\lambda}_{\mathcal{P}}(\Delta_{\Sigma_1}, \Delta_{\Sigma_2}))$, and the other by positive endpoints. Let $\gamma \in \partial_{\infty}\mathbb{H}^3$ be an oriented, piecewise circular curve formed by concatenating the circular arcs bounded by negative endpoints from Δ_{Σ_1} to Δ_{Σ_2} . Let $v_{\Delta_{\Sigma_1}}$ be the

outward tangent vector to the circular arc of the intersection of $\tilde{f}(\Delta_{\Sigma_1}) \cap \partial_\infty \mathbb{H}^3$, and let $v_{\Delta_{\Sigma_2}}$ be the inward tangent vector to the circular arc of the intersection $\tilde{f}(\Delta_{\Sigma_2}) \cap \partial_\infty \mathbb{H}^3$. Then (cf. [3])

$$\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma_2}) = \angle(v_{\Delta_{\Sigma_1}}, v_{\Delta_{\Sigma_2}}) - \sum_W \beta_W$$

where $\angle(v_{\Delta_{\Sigma_1}}, v_{\Delta_{\Sigma_2}})$ is the angle under Euclidean parallel transport in \mathbb{C} , β_W is the signed curvature of the circular subarc W of γ and the sum is over all circular subarcs of γ . In our case, all circular arcs are Euclidean segments and each term of the sum in the above formula is zero. Thus we obtain

$$\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma_2}) = \angle(v_{\Delta_{\Sigma_1}}, v_{\Delta_{\Sigma_2}}).$$

To finish the proof we refer to Figure 3. The vector $v_{\Delta_{\Sigma_1}}$ is parallel to the vector \overrightarrow{xy} in $\partial_\infty \mathbb{H}^3$ whose initial point x is an endpoint of $\tilde{f}(\tilde{C}_1^1)$ and terminal point y is an endpoint of $\tilde{f}(\tilde{C}_2^1)$ as in Figure 3. Similarly, the vector $v_{\Delta_{\Sigma_2}}$ is parallel to the vector $\overrightarrow{y'x'}$ in $\partial_\infty \mathbb{H}^3$ whose initial point y' is an endpoint of $\tilde{f}(\tilde{C}_2^2)$ and terminal point x' is an endpoint of $\tilde{f}(\tilde{C}_1^2)$ as in Figure 3. We normalize the situation such that the short sides h_1 and h_2 of Σ_1 meet \tilde{C} at $j = (0, 0, 1) \in \mathbb{H}^3$ and $e^{-Re(hl(C))}j = Ce^{-R/2}j \in \mathbb{H}^3$ for $e^{-\frac{\epsilon}{R}} < C < e^{\frac{\epsilon}{R}}$. In this case, the points where the short sides h'_1 and h'_2 of Σ_2 meet \tilde{C} are $e^{Re(s(C))}j = Cej \in \mathbb{H}^3$, $e^{-\frac{\epsilon}{R}} < C < e^{\frac{\epsilon}{R}}$, and $e^{Re(s(C)-hl(C))}j = C_1e^{1-\frac{R}{2}}j \in \mathbb{H}^3$, $e^{-\frac{2\epsilon}{R}} < C_1 < e^{\frac{2\epsilon}{R}}$.

If h_1 lies in the xz -plane in \mathbb{H}^3 then x is an analytic function of the complex length $l(h_1)$ of h_1 . An explicit (and elementary) computation shows that the derivative of x in the variable $l(h_1)$ at the point $l(h_1) = 0$ is non-zero. Thus the euclidean distance from x to $0 \in \partial_\infty \mathbb{H}^3$ is $O(|l(h_1)|) = O(e^{-R/4})$ (this holds without the restriction that h_1 is in the xz -plane). Since y is the image of x under the map $z \mapsto e^{-hl(C)}z$, it follows that the distance between y and 0 is $O(e^{-3R/4})$. Similar statements hold for x' and y' , respectively. Consider the Fenchel-Nielsen coordinates $\{(Re(hl(C)), s(C))\}_{C \in \mathcal{P}}$ and let \tilde{f}_{Re} be the corresponding developing map. We normalize \tilde{f}_{Re} such that $\tilde{f}_{Re}(\tilde{C})$ has endpoints 0 and ∞ , and that $\tilde{f}_{Re}(\Delta_{\Sigma_1})$ and $\tilde{f}_{Re}(\Delta_{\Sigma_2})$ have a common endpoint ∞ . Moreover, we require that the common orthogonal between $\tilde{f}_{Re}(\tilde{C})$ and $\tilde{f}_{Re}(\tilde{C}_1^1)$ meets $\tilde{f}_{Re}(\tilde{C})$ at $j \in \mathbb{H}^3$, and the common orthogonal between $\tilde{f}_{Re}(\tilde{C})$ and $\tilde{f}_{Re}(\tilde{C}_1^2)$ meets $\tilde{f}_{Re}(\tilde{C})$ at $e^{1+O(\epsilon/R)}j \in \mathbb{H}^3$. Let x_0, y_0, x'_0 and y'_0 be the endpoints of $\tilde{f}_{Re}(\tilde{C}_1^1)$, $\tilde{f}_{Re}(\tilde{C}_2^1)$, $\tilde{f}_{Re}(\tilde{C}_1^2)$ and $\tilde{f}_{Re}(\tilde{C}_2^2)$ that are different from ∞ , respectively.

Let \tilde{f}_{Im} be the developing map which maps the pleated surface for $\{(Re(hl(C)), s(C))\}_{C \in \mathcal{P}}$ to the pleated surface for $\{(hl(C), s(C))\}_{C \in \mathcal{P}}$ and fixes \tilde{C} . Then $x = \tilde{f}_{Im}(x_0)$, $y = \tilde{f}_{Im}(y_0)$, $x' = \tilde{f}_{Im}(x'_0)$ and $y' = \tilde{f}_{Im}(y'_0)$. In terms of the geometry, x is the image of the endpoint x_0 of the geodesic $\tilde{f}_{Re}(\tilde{C}_1^1)$ under the rotation around the common orthogonal to $\tilde{f}_{Re}(\tilde{C}) = \tilde{C}$ and $\tilde{f}_{Re}(\tilde{C}_1^1)$ with the angle of the rotation equal to the imaginary part of the complex length of the common orthogonal to $\tilde{f}(\tilde{C}) = \tilde{C}$ and $\tilde{f}(\tilde{C}_1^1)$. The cosine formula estimates this angle to be $O(\frac{\epsilon}{R}e^{-R/4})$. Then the euclidean distance between x_0 and x is $O(\frac{\epsilon}{R}e^{-R/4})$ which implies that the distance between x and 0 is $O(e^{-R/4})$. Note that y_0 and y are the images of x_0 and x under the maps $z \mapsto e^{-R/4+(1+i)O(\frac{\epsilon}{R})}$. Thus the euclidean distance between y and y_0 is $O(\frac{\epsilon}{R}e^{-3R/4})$ and the distance between y and 0 is $O(e^{-3R/4})$. Similar properties hold for x'_0, x' and y'_0, y' . The angle between the vectors $\overrightarrow{x_0y_0}$ and $\overrightarrow{y'_0x'_0}$

is $s(C)$ because the length of $C \in \mathcal{P}$ is real (which means that the hexagons are not skewed). The above shows that the angle $\overrightarrow{x_0 y_0}$ and $\overrightarrow{x \tilde{y}}$ is $O(\frac{\epsilon}{R})$, and the same estimate for the angle between $\overrightarrow{y'_0 x'_0}$ and $\overrightarrow{y' x'}$. Thus the angle between $\overrightarrow{x \tilde{y}}$ and $\overrightarrow{y' x'}$ is $O(\frac{\epsilon}{R})$ which finishes the proof. \square

The definition of the bending map. Let Δ_1 and Δ_2 be two complementary triangles to $\tilde{\lambda}$. Bonahon [3] defined the bending map $\tilde{f}_\beta|_{\Delta_2} = \varphi_{\Delta_1, \Delta_2}$ normalized to be the identity at Δ_1 as follows. Let $\mathcal{P}_p = \{\Delta'_1, \Delta'_2, \dots, \Delta'_p\}$ be a sequence of complementary triangles to $\tilde{\lambda}$ which separate Δ_1 and Δ_2 given in the order from Δ_1 to Δ_2 . Define

$$\psi_p = R_{g_{\Delta'_1}^{\Delta_1}}^{\beta(\Delta_1, \Delta'_1)} \circ R_{g_{\Delta'_1}^{\Delta_2}}^{-\beta(\Delta_1, \Delta'_1)} \circ R_{g_{\Delta'_2}^{\Delta_1}}^{\beta(\Delta_1, \Delta'_2)} \circ R_{g_{\Delta'_2}^{\Delta_2}}^{-\beta(\Delta_1, \Delta'_2)} \circ \dots \circ R_{g_{\Delta'_p}^{\Delta_1}}^{\beta(\Delta_1, \Delta'_p)} \circ R_{g_{\Delta'_p}^{\Delta_2}}^{-\beta(\Delta_1, \Delta'_p)}$$

where R_g^b is the hyperbolic rotation around the axis $g \subset \mathbb{H}^3$ by the angle $b \in \mathbb{R}$, and $g_{\Delta'_i}^{\Delta_k}$ is the geodesic on the boundary of Δ'_i which is closest to Δ_k for $k = 1, 2$. Let \mathcal{P} be the family of all complementary triangles to $\tilde{\lambda}$ that separate Δ_1 and Δ_2 . If $\mathcal{P}_p \rightarrow \mathcal{P}$ in the sense that \mathcal{P}_p is an increasing family with $\cup_{p=1}^\infty \mathcal{P}_p = \mathcal{P}$, then the limit

$$\psi_{\Delta_1, \Delta_2} = \lim_{\mathcal{P}_p \rightarrow \mathcal{P}} \psi_p$$

exists and it is independent of the choice of \mathcal{P}_p (cf. [3]). Then

$$\varphi_{\Delta_1, \Delta_2} = \psi_{\Delta_1, \Delta_2} \circ R_{g_{\Delta_2}^{\Delta_1}}^{\beta(\Delta_1, \Delta_2)}.$$

The following lemma is established in [9]. We give a different proof below.

Lemma 3.2. *Under the above assumptions, a geodesic arc in \mathbb{H}^2 of length 1 intersects at most $2R + 2$ geodesics from $\pi^{-1}(\mathcal{P})$, when R is large enough.*

Proof. Let l be an arc of length 1 which transversely intersects geodesics of $\tilde{\mathcal{P}} = \pi^{-1}(\mathcal{P})$. Let $\{\tilde{C}_1, \tilde{C}_2, \dots, \tilde{C}_n\}$ be the geodesics in $\tilde{\mathcal{P}}$ which intersect l in the given order and we orient them to the left as seen from the half-plane in $\mathbb{H}^2 \setminus \tilde{C}_1$ which does not contain \tilde{C}_2 . For R large enough, consecutive geodesics in $\{\tilde{C}_1, \tilde{C}_2, \dots, \tilde{C}_n\}$ are connected by the short arcs of the hexagons (otherwise l would intersect two short sides of a single hexagon which would imply $|l| \geq R/4 > 1$). Given \tilde{C}_j and \tilde{C}_{j+1} , let h_j be the common orthogonal, and let $x_j^+ = \tilde{C}_j \cap h_j$ and $x_{j+1}^- = \tilde{C}_{j+1} \cap h_j$ for $j = 1, \dots, n-1$, and $x_n^+ = x_n^-$. Given $a \in \tilde{C}_j$, define $r(a)$ to be the signed distance between a and x_j^+ . Issue a geodesic g_a through a such that the angle of intersection between \tilde{C}_j and g_a is equal to the angle of intersection between \tilde{C}_{j+1} and g_a . Let $a' = g_a \cap \tilde{C}_{j+1}$. Then the signed distance between a' and x_{j+1}^- is equal to $r(a)$ and consequently the signed distance between a' and x_{j+1}^+ is $r(a) - (1 \pm \frac{\epsilon}{R})$ because the twist parameter is $1 \pm \frac{\epsilon}{R}$ by the assumption. Let $r^-(a)$ denote the signed distance of a to x_j^- for $a \in \tilde{C}_j$. Thus $r^-(a') = r(a)$ for $a \in \tilde{C}_j$.

Let $L_j = l \cap \tilde{C}_j$. We compare the signed distance between L_{j+1} and x_{j+1}^+ to the signed distance between L'_j and x_{j+1}^+ . Consider the hyperbolic triangle with vertices L_j , L'_j and L_{j+1} . The angle at L_j is smaller than the angle at L'_j (cf. Figure 4). By the sine formula for hyperbolic triangles we get

$$(7) \quad d(L_j, L_{j+1}) > d(L'_j, L_{j+1}).$$




FIGURE 4. The number of intersections.

By the condition on the twist parameters we have

$$r(L_{j+1}) = r^-(L_{j+1}) - (1 \pm \frac{\epsilon}{R}).$$

Moreover, we have

$$r^-(L_{j+1}) = r^-(L'_j) \pm d(L'_j, L_{j+1}) = r(L_j) \pm d(L'_j, L_{j+1}).$$

The above gives

$$r(L_{j+1}) = r(L_j) \pm d(L'_j, L_{j+1}) - (1 \pm \epsilon)$$

and thus

$$(8) \quad r(L_n) \geq r(L_1) - 1 + (n-1)(1 \pm \frac{\epsilon}{R})$$

for all $n \geq 2$ because $\sum_{i=1}^n d(L'_i, L_{j+1}) \leq \sum_{i=1}^n d(L_j, L_{j+1}) \leq 1$ by (7).

Assume that $n \geq 2R + 2$. Then there is $1 \leq j \leq n$ such that $|r(L_j)| \geq R$ by (8). We find the contradiction with this inequality by proving that $d(L_j, L_{j+1})$ is too large in this case.

We prove that $d(L_j, L_{j+1})$ is too large. Without the loss of generality, we assume that the quadrilateral Q with vertices L_j, x_j^+, x_{j+1}^- and L_{j+1} has a right angle at the vertex L_{j+1} . It follows then that three angles of Q are equal to $\frac{\pi}{2}$ since the angles at x_j^+ and x_{j+1}^- are equal to $\frac{\pi}{2}$. An elementary hyperbolic geometry gives

$$\cosh^2 d(L_j, L_{j+1}) = \cosh^2 d(x_j^+, L_j) \sinh^2 d(x_j^+, x_{j+1}^-) + 1.$$

Since

$$d(x_j^+, L_j) = |r(L_j)| \geq R$$

and

$$d(x_j^+, x_{j+1}^-) \leq C e^{-R/2},$$

the above gives

$$d(L_j, L_{j+1}) \geq \frac{R}{2} - C$$

for a fixed $C > 0$ and R large enough. This implies that $d(L_j, L_{j+1}) > 1$ for R large enough which is a contradiction. Thus a geodesic arc of length 1 intersects at most $2R + 2$ geodesics of $\tilde{\mathcal{P}}$. \square

4. INJECTIVITY OF THE BENDING MAPS

The purpose of this section is to prove the following theorem which is the first statement of Theorem 1.3 from Introduction. We finish the proof of the remaining statements of Theorem 1.3 in the next section.

Theorem 4.1. *Given $C_0 > 0$, there exist $\hat{\epsilon} > 0$ and $R(\hat{\epsilon}) > 0$ such that for each $0 \leq \epsilon < \hat{\epsilon}$ and $R \geq R(\hat{\epsilon})$ the following is satisfied. Let S be a closed hyperbolic surface equipped with a maximal, finite geodesic lamination λ such that each closed geodesic of λ has length in the interval $(R - \frac{\epsilon}{R}, R + \frac{\epsilon}{R})$ and that each geodesic arc of length 1 intersects at most $C_0 R$ closed geodesics of λ . Assume that a bending cocycle β transverse to the lift $\tilde{\lambda}$ in \mathbb{H}^2 satisfies*

$$(9) \quad |\beta(\Delta^1(\tilde{l}), \Delta^2(\tilde{l}))| \leq \frac{C_0 \epsilon}{R}$$

for each isolated leaf \tilde{l} and complementary triangles $\Delta^1(\tilde{l})$ and $\Delta^2(\tilde{l})$ with common boundary \tilde{l} , and

$$(10) \quad |\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma_2})| \leq \frac{C_0 \epsilon}{R}$$

for the characteristic triangles Δ_{Σ_1} and Δ_{Σ_2} of each two 0-neighbors hexagons Σ_1 and Σ_2 (coming from the pants decomposition of S whose cuffs are closed geodesics of λ). Then the induced bending map

$$\tilde{f}_\beta : \partial_\infty \mathbb{H}^2 \rightarrow \partial_\infty \mathbb{H}^3$$

is injective.

Proof. Let $x, y \in \partial_\infty \mathbb{H}^2$ be two different points. We need to prove that $\tilde{f}_\beta(x) \neq \tilde{f}_\beta(y)$. Let \mathcal{P} be a pants decomposition of S whose cuffs are closed curves of λ . We fix a decomposition of S into hexagons as in §3 using the pants decomposition \mathcal{P} and lift it to the universal covering $\pi : \mathbb{H}^2 \rightarrow S$. Recall that $\tilde{\lambda} = \pi^{-1}(\lambda)$ and $\tilde{\mathcal{P}} = \pi^{-1}(\mathcal{P})$. Let g be the geodesic in \mathbb{H}^2 whose ideal endpoints are x and y . If $g \subset \mathbb{H}^2 \setminus \mathcal{TH}_t$ then g is a lift of some $C \in \mathcal{P}$ and $\tilde{f}_\beta(x) \neq \tilde{f}_\beta(y)$ because g is in the bending locus of \tilde{f}_β .

Therefore we assume that $g \cap \mathcal{TH}_t \neq \emptyset$. Fix a hexagon Σ^0 such that $g \cap \Sigma_t^0 \neq \emptyset$. Let P be a point in $g \cap \Sigma_t^0$. The point P divides the geodesic g into two rays $g_{\pm 1}$. Let $P_0 = P$ and assume that we have chosen points $P_{\pm 1}, P_{\pm 2}, \dots, P_{\pm n}$ in the increasing order on $g_{\pm 1}$ such that $P_{\pm k} \in (\Sigma_{\pm k})_t$ for distinct hexagons $\Sigma_{\pm k}$, for $k = 1, 2, \dots, n$. We define $P_{\pm(n+1)}$ as follows. Let $\Sigma_{\pm(n+1)}$ be the first hexagon after $\Sigma_{\pm n}$ such that $(\Sigma_{\pm(n+1)})_t$ intersect $g_{\pm 1}$ and that there exists a point $P_{\pm(n+1)} \in g_{\pm 1} \cap (\Sigma_{\pm(n+1)})_t$ with $d(P_{\pm n}, P_{\pm(n+1)}) \geq 1$. If such hexagon does not exist, then we set $P_{\pm(n+1)}$ to be the ideal endpoint of $g_{\pm 1}$. In this fashion we partition each $g_{\pm 1}$ into consecutive arcs of lengths at least 1. It is possible that the partition is finite when $P_{\pm(n+1)}$ is the endpoint of $g_{\pm 1}$.

Let $g \subset \mathbb{H}^3$ be a geodesic ray with initial point p_0 , and let $p \in g$ be another point. For $0 < \theta < \pi$, the cone $\mathcal{C}(p, g, \theta)$ with vertex p , axis g and angle θ is the set of all $w \in \mathbb{H}^3$ such that the angle at p between the positive direction of g and the geodesic ray from p through w is less than θ . Note that a cone is an open set. A non-zero vector $(p, v) \in T^1(\mathbb{H}^3)$ uniquely determines a geodesic ray g which starts at the basepoint p of v and which is tangent to v . Then $\mathcal{C}(p, v, \theta)$ is by the definition $\mathcal{C}(p, g, \theta)$. The shadow of the cone $\mathcal{C}(p, g, \theta)$ is the set $\partial_\infty \mathcal{C}(p, g, \theta)$ of endpoints at

$\partial_\infty \mathbb{H}^3$ of all geodesic rays starting at p and inside $\mathcal{C}(p, g, \theta)$. The shadow of a cone is an open subset of $\partial_\infty \mathbb{H}^3$.

For $d > 0$, let $p_d \in g$ be the point on g which is on the distance d from $p_0 = p$. Let $\eta > 0$ be the maximal angle such that $\mathcal{C}(p_d, g, \eta) \subset \mathcal{C}(p_0, g, \theta)$. Then $\eta = \eta(d, \theta)$ is a continuous function of d and θ . For a fixed $0 < \theta < \pi$, we have $\eta(d, \theta) > \theta$ and $\eta(d, \theta) \rightarrow \theta$ as $d \rightarrow 0$. These properties are elementary.

Let $\{P_{\pm n}\}_n$ be the points of the partition of $g_{\pm 1}$. We consider a sequence of cones $\{\mathcal{C}(P_{\pm n}, g_{\pm 1}, \frac{\pi}{2})\}$. Then

$$\overline{\partial_\infty(\mathcal{C}(P_{\pm(n+1)}, g_{\pm 1}, \frac{\pi}{2}))} \subset \partial_\infty(\mathcal{C}(P_{\pm n}, g_{\pm 1}, \frac{\pi}{2}))$$

for each $n \in \mathbb{N}$ and we say that the sequence of cones is *nested*.

If we prove that the images of the nested cones under the bending map \tilde{f}_β remain nested then we are done. Indeed, since x and y lie in the intersection of the shadows of all nested cones along g_1 and g_{-1} , since the shadows of $\mathcal{C}(P_0, g_1, \frac{\pi}{2})$ and $\mathcal{C}(P_0, g_{-1}, \frac{\pi}{2})$ are disjoint, and if \tilde{f}_β preserves the nesting of the cones, it follows that $\tilde{f}_\beta(x) \neq \tilde{f}_\beta(y)$. It remains to prove that \tilde{f}_β preserves the nesting of the cones. To see this, it is enough to normalize \tilde{f}_β to be the identity on the canonical triangle $\Delta_{\Sigma_{\pm n}}$ of $\Sigma_{\pm n}$ and to prove that

$$\overline{(\tilde{f}_\beta|_{\Delta_{\Sigma_{\pm(n+1)}}})(\partial_\infty \mathcal{C}(P_{\pm(n+1)}, g_{\pm 1}, \frac{\pi}{2}))} \subset \partial_\infty \mathcal{C}(P_{\pm n}, g_{\pm 1}, \frac{\pi}{2})$$

for each $n \in \mathbb{N}$.

Let $a_{\pm n}$ be the arc of $g_{\pm 1}$ between $P_{\pm n}$ and $P_{\pm(n+1)}$. Note that the length of $a_{\pm n}$ is at least 1 and that it can be infinite. We first assume that $a_{\pm n}$ has finite length. Let $\Sigma_1, \Sigma_2, \dots, \Sigma_k$ be the sequence of all hexagons such that $(\Sigma_i)_t \cap a_{\pm n} \neq \emptyset$, for $i = 1, 2, \dots, k$. Note that $(\Sigma_1)_t \ni P_{\pm n}$ and $(\Sigma_k)_t \ni P_{\pm(n+1)}$. For a hexagon Σ , define $C(\Sigma)$ to be the union of all hexagons which are connected by a sequence of 0-neighbors to Σ . Note that $C(\Sigma)$ looks like a trivalent tree and that it has infinitely many boundary components which are made out of partial boundaries of the hexagons in $C(\Sigma)$. It is important to note that either $C(\Sigma_1) = C(\Sigma_{k-1})$, or $C(\Sigma_1)$ and $C(\Sigma_{k-1})$ share a boundary component. If not, then the subarc of $a_{\pm n}$ which connects $(\Sigma_1)_t$ to $(\Sigma_{k-1})_t$ connects two boundary components of some $C(\Sigma')$, where $C(\Sigma')$ separates $C(\Sigma_1)$ and $C(\Sigma_{k-1})$. Note that the arc which connects a short side of a hexagon to a non-adjacent side of the same hexagon has length at least $R/4$, where the long sides of the hexagon have lengths $R/2$. It follows that the subarc of $a_{\pm n}$ which connects two boundary components of $C(\Sigma')$ has length at least $R/4 - 3$. Thus the above subarc of $a_{\pm n}$ has length greater than 1 when R is large enough which is impossible.

If $C(\Sigma_1) = C(\Sigma_{k-1})$ then we form a new sequence of hexagons $\Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-2}, \Sigma_{k-1}$ such that the adjacent pairs of hexagons are 0-neighbors and $a_{\pm n}$ intersects characteristic triangles of the hexagons in the sequence. If $C(\Sigma_1) \neq C(\Sigma_{k-1})$ (and they share a boundary component) then we can choose a new sequence of hexagons $\Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-2}, \Sigma_{k-1}$ such that each pair of adjacent hexagons are 0-neighbors except one adjacent pair that are 1-neighbors, and that $a_{\pm n}$ intersects characteristic triangles of the sequence. Note that the subarc of $a_{\pm n}$ that connects $(\Sigma_1)_t$ and $(\Sigma_{k-1})_t$ is of length less than 1.

The hexagons Σ_{k-1} and Σ_k are either 0- or 1-neighbors, or neither 0- nor 1-neighbors. If Σ_{k-1} and Σ_k are either 0- or 1-neighbors, then $\Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-2}, \Sigma_{k-1}, \Sigma_k$

is a sequence of hexagons whose adjacent hexagons are 0-neighbors with the exception of at most 2 pairs which are 1-neighbors. Note that the arc $a_{\pm n}$ could have large length in general. If Σ_k is a 0-neighbor of Σ_{k-1} then there is an arc $b_{\pm n}$ from the second point of the intersection of $a_{\pm n}$ with the boundary of $(\Sigma_{k-1})_t$ to the boundary of $(\Sigma_k)_t$ that has length less than 2. To see this, let $\tilde{C} \in \pi^{-1}(\mathcal{P}) = \tilde{\mathcal{P}}$ be the geodesic which contains one boundary side of both Σ_{k-1} and Σ_k . Then the boundary side of $(\Sigma_{k-1})_t$ closest to \tilde{C} is in the $C_1 e^{-R/4}$ -neighborhood of \tilde{C} for some $C_1 > 0$, and the same statement is true for $(\Sigma_k)_t$. Since Σ_k is shifted by $1 \pm \frac{\epsilon}{R}$ with respect to Σ_{k-1} , it follows that such $b_{\pm n}$ exists. Thus the set of geodesics of $\tilde{\lambda} = \pi^{-1}(\lambda)$ that intersect $a_{\pm n}$ also intersect a geodesic arc $c_{\pm n}$ with the initial point $P_{\pm n}$ and of length at most 3. Assume now that Σ_k and Σ_{k-1} are 1-neighbors and that $\tilde{C} \in \tilde{\mathcal{P}}$ separates them. Let Σ'_k be the 0-neighbor of Σ_{k-1} which is separated by \tilde{C} from Σ_{k-1} . It follows that the geodesics of $\tilde{\lambda}$ which intersect $a_{\pm n}$ except possibly the last geodesic (namely, the geodesic which contains one side of $(\Sigma_k)_t$ closest to \tilde{C}) intersect a geodesic arc of length at most 3 with one endpoint $P_{\pm n}$. This follows simply by applying the above reasoning to the sequence $\Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-2}, \Sigma_{k-1}, \Sigma'_k$. If Σ_k and Σ'_k are not separated by some $\tilde{C} \in \mathcal{P}$, then $\Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-2}, \Sigma_{k-1}$ suffices to get the same conclusion.

We give a proof of the nesting for the second case discussed above and the first case above is a subcase of the second. Namely, we are assuming that the set of geodesics $\tilde{\lambda}(a_{\pm n})$ of $\tilde{\lambda}$ which intersect $a_{\pm n}$ is also intersected by a geodesic arc $c_{\pm n}$ of length at most 3 with the initial point $P_{\pm n}$ with a possible exception of one geodesic in $\tilde{\lambda}(a_{\pm n})$. We consider the bending map $(\tilde{f}_\beta)|_{\Delta_{\Sigma_k}} = \varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma_k}}$. Let g_k be the geodesic of $\tilde{\lambda}$ which contains the boundary of $(\Sigma_k)_t$ and that separates $(\Sigma_k)_t$ and $(\Sigma'_k)_t$. If $(\Sigma_k)_t \cap a_{\pm n}$ comes before $(\Sigma'_k)_t \cap a_{\pm n}$ along $a_{\pm n}$ then $c_{\pm n}$ does intersect Δ_{Σ_k} and this subcase of the second case reduces to the first case. Therefore, we assume that $(\Sigma_k)_t \cap a_{\pm n}$ comes after $(\Sigma'_k)_t \cap a_{\pm n}$ along $a_{\pm n}$. The geodesic g_k might not intersect $c_{\pm n}$. We have

$$\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma_k}} = \varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}} \circ R_{g_k}^{\beta(g_k)}$$

where Σ'_k is the 0-neighbor of Σ_{k-1} that is separated from Σ_k by the geodesic g_k .

We normalize such that $P_{\pm n} = j \in \mathbb{H}^3$ and $P_{\pm(n+1)} = e^{-m}j$, where $m \geq 1$. Then $v = \{e^{-m}j, -j\}$ is a tangent vector to $a_{\pm n}$ at the point $P_{\pm(n+1)}$ pointing towards the ideal endpoint of $g_{\pm 1}$. Lemma A.3 and the assumptions give

$$D_{T\mathbb{H}^3}(R_{g_k}^{\beta(\Delta_1(g_k), \Delta_2(g_k))})(\{e^{-m}j, -j\}), \{e^{-m}j, -j\}) \leq C|\beta(\Delta_1(g_k), \Delta_2(g_k))| \leq \frac{C'\epsilon}{R} \quad (11)$$

for some $C' > 0$ when $\epsilon > 0$ is small enough and $R \geq 1$, where $\{e^{-m}j, -j\} \in T\mathbb{H}^3$ is a tangent vector to \mathbb{H}^3 based at $e^{-m}j$ and the function $D_{T\mathbb{H}^3}(\cdot, \cdot)$ is defined in Appendix formula (14).

We consider

$$\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}} = \psi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}} \circ R_{g'_k}^{\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma'_k})}$$

where $g'_k \in \tilde{\lambda}$ is the side of $\Delta_{\Sigma'_k}$ facing Δ_{Σ_1} . Let $\Delta_{\Sigma'_i}$ and $\Delta_{\Sigma'_{i+1}}$ be canonical triangles of two adjacent hexagons from the sequence $\Sigma'_1 := \Sigma_1, \Sigma'_2, \dots, \Sigma'_{k-1} := \Sigma_{k-1}, \Sigma'_k$. Then $\Delta_{\Sigma'_i}$ and $\Delta_{\Sigma'_{i+1}}$ are separated by $\tilde{C}_i \in \tilde{\mathcal{P}}$ and they have a common endpoint \tilde{x}_i with \tilde{C}_i . Let \mathcal{F}_i be the family of complementary triangles between $\Delta_{\Sigma'_i}$




FIGURE 5.

and $\Delta_{\Sigma'_{i+1}}$. Let Δ_1^i, Δ_2^i be two triangles in \mathcal{F}_i which are closets to $\Delta_{\Sigma'_i}$ and let $\Delta_1^{i+1}, \Delta_2^{i+1}$ be two triangles in \mathcal{F}_i which are closets to $\Delta_{\Sigma'_{i+1}}$. Let $\gamma_i \in PSL_2(\mathbb{R})$ be the element of the covering group of S that corresponds to \tilde{C}_i . Any triangle in \mathcal{F}_i between $\Delta_{\Sigma'_i}$ and \tilde{C}_i is the image of either Δ_1^i or Δ_2^i under a power of γ_i , and any triangle of \mathcal{F}_i between \tilde{C}_i and $\Delta_{\Sigma'_{i+1}}$ is the image of either Δ_1^{i+1} or Δ_2^{i+1} under a power of γ_i .

Let h_i be the horocyclic arc connecting $\Delta_{\Sigma'_i}$ and $\Delta_{\Sigma'_{i+1}}$ with the center $\tilde{y}_i = c_{\pm n} \cap \tilde{C}_i$. Note that the length $|h_i|$ of the arc h_i is less than a constant multiple of the length of the subarc of $c_{\pm n}$ connecting $\Delta_{\Sigma'_i}$ and $\Delta_{\Sigma'_{i+1}}$. Moreover, the length $|h_i|$ is less than the sum of the lengths of $h_i \cap \Delta_k^i, h_i \cap \Delta_k^{i+1}$, for $k = 1, 2$, and of the sum of the lengths of the intersections of h_i with the translates of Δ_k^i and Δ_k^{i+1} , for $k = 1, 2$, under the powers of γ_i such that the common endpoint of Δ_1^i and \tilde{C}_i is repelling. Then the length of all translates is less than $C|h_i|e^{-l(\gamma_i)/2}$, where $l(\gamma_i)$ is the real part of the translation length of γ_i . To see this, we normalize the situation such that $\tilde{y}_i = (0, 1) \in \mathbb{H}^2$ and \tilde{C}_i is the geodesic with endpoints 0 and ∞ . Then h_i is the horizontal Euclidean arc which contains $(0, 1) \in \mathbb{H}^2$ and the Euclidean length of h_i equals the hyperbolic length of h_i . Let $\{f_m\}_{m \in \mathbb{N}}$ be the geodesics of $\tilde{\lambda}$ with one endpoint ∞ that separate $\Delta_{\Sigma'_i}$ and \tilde{C}_i in the increasing order from $\Delta_{\Sigma'_i}$. Let $\{e_m\}_{m \in \mathbb{N}}$ be the geodesics of $\tilde{\lambda}$ with one endpoint ∞ that separate $\Delta_{\Sigma'_{i+1}}$ and \tilde{C}_i in the decreasing order from $\Delta_{\Sigma'_{i+1}}$. Let x_1 be the length of the arc of h_i between f_1 and f_2 , and let x_2 be the length of the arc of h_i between f_2 and f_3 . Note that f_m is mapped to f_{m+2} by the hyperbolic translation γ_i with the axis \tilde{C}_i and the attracting fixed point $0 \in \partial_\infty \mathbb{H}^2$. Then the distance between f_{2m+1} and f_{2m+2} is $x_1 e^{-\frac{ml(\gamma_i)}{2}}$, and similarly the distance between f_{2m+2} and f_{2m+3} is $x_2 e^{-\frac{ml(\gamma_i)}{2}}$ for $m \in \mathbb{N}$ (cf. Figure 5). Therefore, the sum of the lengths of the gaps of h_i except the first two gaps and the last two gaps is bounded by $C|h_i|e^{-l(\gamma_i)/2}$.

Note that

$$(12) \quad \begin{aligned} |\beta(\Delta_{\Sigma'_i}, \Delta_1^i)| &\leq \frac{C_0\epsilon}{R} \\ |\beta(\Delta_{\Sigma'_i}, \Delta_2^i)| &\leq \frac{2C_0\epsilon}{R} \\ |\beta(\Delta_{\Sigma'_i}, \Delta_{\Sigma'_{i+1}})| &\leq \frac{C_0\epsilon}{R} \\ |\beta(\Delta_{\Sigma'_i}, \Delta_1^{i+1})| &\leq \frac{2C_0\epsilon}{R} \\ |\beta(\Delta_{\Sigma'_i}, \Delta_2^{i+1})| &\leq \frac{3C_0\epsilon}{R}. \end{aligned}$$

By the uniform boundedness of the composition of rotations [3], there exists $C > 0$ such that

$$\|\psi_{\Delta_{\Sigma_1}, \Delta_{\Sigma_k}} - id\| \leq C \sum_d \|R_{g_d^{\Delta_{\Sigma_1}}}^{\beta(k_d)} R_{g_d^{\Delta_{\Sigma_k}}}^{-\beta(k_d)} - id\|$$

where the sum is over all gaps d of $c_{\pm n}$, k_d is the subarc of $c_{\pm n}$ from $P_{\pm n}$ to a point in d , and $g_d^{\Delta_{\Sigma_1}}$ ($g_d^{\Delta_{\Sigma_k}}$) is the leaf of $\tilde{\lambda}$ which contains the endpoint of d closer to Δ_{Σ_1} (Δ_{Σ_k}). We divide the above sum over the gaps of $c_{\pm n}$ into two sums Σ' and Σ'' . The first sum Σ' is over all gaps $c_{\pm n} \cap \Delta_{\Sigma'_i}$, for $i = 1, 2, \dots, k$, and $c_{\pm n} \cap \Delta_l^i$, for $l = 1, 2$, and the second sum Σ'' is over the remaining gaps.

The first sum is finite. By Lemma 3.2, $k \leq 2R + 2 \leq 4R$ for $R \geq 1$ and by the finite additivity of β , we have that

$$|\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma'_i})| \leq C_1\epsilon$$

for $i = 1, 2, \dots, k$, as well as

$$|\beta(\Delta_{\Sigma_1}, \Delta_l^i)| \leq C_1\epsilon$$

for $i = 1, 2, \dots, k$ and $l = 1, 2$, and some constant $C_1 > 0$.

Lemma A.4 implies

$$\sum' \leq \sum_{i=1}^k C_2\epsilon|h_i| \leq C_2|c_{\pm n}|\epsilon \leq C_3\epsilon.$$

It remains to estimate Σ'' . We proved above that the total length of the gaps of h_i with respect to the family \mathcal{F}_i of complementary triangles except for the first two and the last two gaps is less than $Ce^{-R/2}|h_i|$. Since $\sum_{i=1}^n |h_i| \leq C|c_{\pm n}| \leq C_4$ and the cocycle β takes values in $[-\pi, \pi)$ (thus β is bounded), Lemma A.4 gives

$$\sum'' \leq C \sum_{i=1}^k e^{-R/2} \leq C_5 R e^{-R/2}$$

for some $C_5 > 0$. Then $\Sigma' + \Sigma''$ can be made arbitrary small when $\epsilon > 0$ is small enough and $R > 0$ is large enough. The above, Lemma A.3 and

$$|\beta(\Delta_{\Sigma_1}, \Delta_{\Sigma'_k})| \leq C\epsilon$$

imply that

$$\|\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}} - id\|$$

is as small as needed for $\epsilon > 0$ small enough and $R > 0$ large enough. Then (12), Lemma A.2 and the above prove that the assumptions of Lemma A.1 are satisfied for $\epsilon > 0$ small enough and $R > 0$ large enough. Thus the nesting for \tilde{f}_β on $a_{\pm n}$ follows by Lemma A.1. We choose $\hat{\epsilon} > 0$ and $R(\hat{\epsilon}) > 0$ accordingly.

We assume now that Σ'_{k-1} and Σ_k are neither 0- nor 1-neighbors. Then there is a unique $\tilde{C}_{k-1} \in \tilde{\mathcal{P}}$ which separates Σ'_{k-1} and Σ_k , and that contains boundary

sides of both of them. Let Σ'_k be the 0-neighbor of Σ'_{k-1} which is on the same side of \tilde{C}_{k-1} as Σ_k . Let $s \geq 1$ be the number of hexagons in between Σ'_k and Σ_k . There are two possibilities: either $a_{\pm n}$ intersect $\Delta_{\Sigma'_k}$ in which case we say that Δ_{Σ_k} is “above” $\Delta_{\Sigma'_k}$, or $a_{\pm n}$ does not intersect $\Delta_{\Sigma'_k}$ in which case we say that Δ_{Σ_k} is “below” $\Delta_{\Sigma'_{k-1}}$.

Assume we are in the former case and let $\{g_1, g_2, \dots, g_{s+1}\}$ be the geodesic of $\tilde{\lambda}$ between $\Delta_{\Sigma'_k}$ and Δ_{Σ_k} . We use the following fact. Let h and h' be two geodesics that intersect $L = \{(0, 0, t) : t > 0\}$ at points $e^{-m}j$ and $e^{-m'}j$ subtending angles $\epsilon > 0$ and $\epsilon' > 0$, where $m < m'$. Let $\epsilon'' = \max\{\epsilon, \epsilon'\}$ and let h'' be the geodesic that intersects L at the point $e^{-m}j$ subtending an angle ϵ'' . Then, for $m'' \geq m'$ and $\epsilon'' > 0$, we have

$$\begin{aligned} D_{T\mathbb{H}^3}(R_h^\theta \circ R_{h'}^{\theta'}(\{e^{-m''}j, -j\}), \{e^{-m''}j, -j\}) &\leq \\ &\leq \max_{0 \leq \theta'' \leq 2\pi} D_{T\mathbb{H}^3}(R_{h''}^{\theta''}(\{e^{-m''}j, -j\}), \{e^{-m''}j, -j\}). \end{aligned}$$

Let

$$R_s = R_{g_1}^{\beta(\Delta_1(g_1), \Delta_2(g_1))} \circ \dots \circ R_{g_{s+1}}^{\beta(\Delta_1(g_{s+1}), \Delta_2(g_{s+1}))}.$$

Then, the above implies that

$$(13) \quad \begin{aligned} D_{T\mathbb{H}^3}(R_s(\{e^{-m''}j, -j\}), \{e^{-m''}j, -j\}) &\leq \\ &\leq \max_{0 \leq \theta \leq 2\pi} D_{T\mathbb{H}^3}(R_{g'_1}^\theta(\{e^{-m''}j, -j\}), \{e^{-m''}j, -j\}). \end{aligned}$$

where g'_1 is the geodesic passing through $g_1 \cap L$ that subtends an angle $\max\{|\angle(g_1, L)|, \dots, |\angle(g_s, L)|\}$ with L . Lemma A.5 and (13) imply that $D_{T\mathbb{H}^3}(R_s(\{e^{-m''}j, -j\}), \{e^{-m''}j, -j\})$ is as small as we want when the angle $|\angle(g'_1, L)|$ is small enough for any $0 \leq \theta \leq 2\pi$. Note that

$$\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma_k}} = \varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}} \circ \varphi_{\Delta_{\Sigma'_k}, \Delta_{\Sigma_k}}.$$

Since $\varphi_{\Delta_{\Sigma'_k}, \Delta_{\Sigma_k}} = R_s$, the above gives

$$D_{T\mathbb{H}^3}(\varphi_{\Delta_{\Sigma'_k}, \Delta_{\Sigma_k}}(\{P_{\pm(n+1)}, -j\}), \{P_{\pm(n+1)}, -j\})$$

is as small as needed for R large enough. Indeed, the subarc of $a_{\pm n}$ from the second point of the intersection with the boundary of $(\Sigma_{k-1})_t$ to the first point of intersection with the boundary of $(\Sigma'_k)_t$ is inside one complement of \mathcal{TH}_t as well as long sub arcs of the set of geodesics $\{g_1, \dots, g_{s+1}\}$. Thus $a_{\pm n}$ and $\{g_1, \dots, g_{s+1}\}$ remain in a neighborhood of one $\tilde{C} \in \tilde{\mathcal{P}}$ for a long distance when R is large. It follows that the angles of intersections between $a_{\pm n}$ and the geodesics in $\{g_1, g_2, \dots, g_{s+1}\}$ are small for R large enough and the above applies. The reasoning in the first case applies to $\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma'_k}}$ and we have the nesting of the images of the cones at the endpoints of $a_{\pm n}$ under the bending map $\varphi_{\Delta_{\Sigma_1}, \Delta_{\Sigma_k}}$. If $\Delta_{\Sigma'_{k-1}}$ is “above” Δ_{Σ_k} then symmetry reduces to the previous case.

It remains to consider the case when $a_{\pm n}$ has infinite length (in which case the endpoint of $a_{\pm n}$ is also the endpoint of $\tilde{C} \in \tilde{\mathcal{P}}$ and $a_{\pm n} \subset \mathbb{H}^2 - \mathcal{TH}_t$). An elementary (euclidean) considerations prove that when $R \geq 1$ the number of geodesics of $\tilde{\lambda}$ that intersect the geodesics subrays of $a_{\pm n}$ which connect two sides of a single hexagon is at most 6. Indeed, assume that $a_{\pm n}$ is the geodesic arc in \mathbb{H}^2 with the initial point $i - e^{-R/4}$ and the endpoint $0 \in \partial_\infty \mathbb{H}^2$. Then $a_{\pm n}$ is a circular arc with the center $a = \frac{e^{R/4} + e^{-R/4} - \sqrt{(e^{R/4} + e^{-R/4})^2 - 4e^{-R}}}{2}$ and the radius a . The x -coordinate of

the intersection of $a_{\pm n}$ with the horizontal line $y = e^{-R/2}$ is estimated to be more than $e^{-3R/2}$. Since the translation length of the element γ fixing the y -axis is $e^{-R/2}$ and since γ identifies every second geodesic of $\tilde{\lambda}$ that have endpoint ∞ , the claim follows. Then applying Lemma A.3 finitely many times to the sequence of subarcs of $a_{\pm n}$ of lengths $R/2$, we obtain a nesting property along this sequence. Thus \tilde{f}_β is injective for $\epsilon > 0$ small enough and $R > 0$ large enough.

We choose $\hat{\epsilon}$ and $R(\hat{\epsilon})$ as the minimum of the choice in all the cases considered. \square

5. HOLOMORPHIC MOTIONS

We finish the proof of Theorem 1.3 using holomorphic motions. This proof is standard once the injectivity is established (cf. [8], [7] and [13]). Holomorphic motions were introduced and studied in [11] and the key extension property is proved in [15].

The endpoints of the representations of elements of $\pi_1(S)$ vary holomorphically in the complex Fenchel-Nielsen coordinates. We established that the holomorphic variation is injective on the set of endpoints when the parameters are close to being real in the sense of (2) and (3). Thus the holomorphic variation of the endpoint of $\pi_1(S)$ is a holomorphic motion which extends by the lambda lemma (cf. [11]) to a holomorphic motion of the unit circle. Then there exists an extension to a holomorphic motion of the complex plane (cf. [15]). It follows that \tilde{f}_β extends to a quasiconformal mapping of the complex plane and that the quasiconformal constant is less than $1 + K_0\epsilon$ for $\epsilon > 0$ small enough and fixed $K_0 > 0$ (cf. [11]). The extension of \tilde{f}_β can be chosen to be equivariant with respect to the action of $\pi_1(S)$ (cf. [5]) which finishes the proof of Theorem 1.3.

APPENDIX

We use the quaternions to represent the upper half-space model $\mathbb{H}^3 = \{z + tj : z \in \mathbb{C}, t > 0\}$ of the hyperbolic three-space (see Beardon [1]), where $j = (0, 0, 1) \in \mathbb{H}^3$. The space of isometries of \mathbb{H}^3 is identified with $PSL_2(\mathbb{C})$. The Poincaré extension of $A(z) = \frac{az+b}{cz+d} \in PSL_2(\mathbb{C})$ to \mathbb{H}^3 is given in [1] by

$$A(z + tj) = \frac{(az + b)\overline{(cz + d)} + a\bar{c}t^2 + tj}{|cz + d|^2 + |c|^2t^2}.$$

An isometry of \mathbb{H}^3 which is close to the identity moves points on a bounded distance from $j \in \mathbb{H}^3$ by a small amount and the tangent vectors are rotated by a small angle with respect to the Euclidean parallel transport in \mathbb{R}^3 . We give a quantitative statement for the above including the situation when the points are on the unbounded distances from $j \in \mathbb{H}^3$ which is needed in our considerations.

Given $P = z + tj \in \mathbb{H}^3$, we define

$$ht(P) = t$$

and

$$Z(P) = z.$$

Consider the tangent space $T\mathbb{H}^3$ to the upper half-space \mathbb{H}^3 . Let $\{P, u\}, \{Q, v\} \in T\mathbb{H}^3$ be two tangent vectors based at $P, Q \in \mathbb{H}^3$, respectively. We define

$$(14) \quad D_{T\mathbb{H}^3}(\{P, u\}, \{Q, v\}) = \max\left\{\left|\frac{ht(P)}{ht(Q)} - 1\right|, |Z(P) - Z(Q)|, |\angle(u, v)|\right\},$$




FIGURE 6.

where $\angle(u, v)$ is the angle between the vectors u and v after the euclidean transport in \mathbb{H}^3 . Note that $D_{T\mathbb{H}^3}(\{P, u\}, \{Q, v\})$ is not a metric on $T\mathbb{H}^3$.

Lemma A.1. *Given $m_0 > 0$, there exists $\delta = \delta(m_0) > 0$ such that for any $m \geq m_0$ we have*

$$(15) \quad \overline{\partial_\infty \mathcal{C}(P, v, \frac{\pi}{2})} \subset \partial_\infty \mathcal{C}(j, -j, \frac{\pi}{2})$$

where $\{P, v\} \in T\mathbb{H}^3$ satisfies $D_{T\mathbb{H}^3}(\{P, v\}, \{e^{-m}j, -j\}) < \delta$.

Proof. Let x be the center of the Euclidean hemisphere (orthogonal to \mathbb{C}) that passes through P and touches the unit Euclidean hemisphere orthogonal to \mathbb{C} with the center $0 \in \mathbb{C} \subset \partial_\infty \mathbb{H}^3$ (cf. Figure 6). Let $y = Z(P) \in \mathbb{C}$ and let φ be the angle between euclidean segments Px and Py at the point P .

An elementary (Euclidean) considerations give

$$(16) \quad x \geq C_1(m_0) > 0$$

for some constant $C_1(m_0) > 0$ which depends on m_0 .

This implies that

$$\varphi \geq C_2(m_0) > 0$$

for some constant $C_2(m_0) > 0$. Thus

$$(17) \quad \overline{\partial_\infty \mathcal{C}(P, -j, \frac{\pi}{2} + C_2(m_0))} \subset \partial_\infty \mathcal{C}(j, -j, \frac{\pi}{2}).$$

Since the angle (at the point P) between the hyperbolic geodesic connecting j to P and the euclidean segment Py is less than φ , the above inclusion implies (15) for $\delta(m_0) < C_2(m_0)$. \square

Lemma A.2. *Let $g \in PSL_2(\mathbb{C})$ with*

$$\|g - id\| < \frac{1}{2}$$

and let $\{z + tj, u\}$ be a tangent vector to \mathbb{H}^3 such that

$$D_{T\mathbb{H}^3}(\{z + tj, u\}, \{e^{-m}j, -j\}) \leq \delta$$

for $0 \leq \delta < \delta_0$ with $\delta_0 > 0$ fixed. Then there exist $C_1, C_2 > 0$ depending on δ_0 such that for every $m > 0$ we have

$$D_{T\mathbb{H}^3}(g(\{z + tj, u\}), \{e^{-m}j, -j\}) \leq C_1\delta + C_2\|g - id\|.$$

Proof. Denote by g the Poincaré extension of $g(z) = \frac{az+b}{cz+d}$ with $ad - bc = 1$ given above. Then

$$\left|ht(g(z + tj)) - e^{-m}\right| = \left|\frac{t}{|cz + d|^2 + |c|^2t^2} - e^{-m}\right| \leq C_1e^{-m}\delta + C_2e^{-m}\|g - id\|$$

for constants $C_1, C_2 > 0$ independent of $m > 0$ and for all $g \in PSL_2(\mathbf{C})$ with $\|g - id\| \leq \frac{1}{2}$. Moreover,

$$\left|Z(g(z + tj))\right| = \left|\frac{a\bar{c}|z|^2 + a\bar{d}z + b\bar{c}\bar{z} + b\bar{d} + a\bar{c}t^2}{|cz + d|^2 + |c|^2t^2}\right| \leq C_1\delta + C_2\|g - id\|.$$

Let $u = \langle u_1, u_2, u_3 \rangle$. Without loss of generality, we assume that $|u_1|, |u_2| \leq \delta$ and $u_3 = -j$. Let $v = Dg(z + tj)u = \langle v_1, v_2, v_3 \rangle$. Let $g = g_1 + g_2i + g_3j$ be the coordinate functions of g . Direct computations give

$$\left|\frac{\partial g_i}{\partial x}(z + tj)\right|, \left|\frac{\partial g_i}{\partial y}(z + tj)\right| \leq C$$

for some $C > 0$ and $i = 1, 2, 3$, where $z + tj = x + yi + tj$. Moreover,

$$\left|\frac{\partial g_i}{\partial t}(z + tj)\right| \leq C_1\|g - id\|$$

for some $C_1 > 0$ and $i = 1, 2$, and

$$\left|\frac{\partial g_3}{\partial t}(z + tj)\right| \geq 1 - C_2\|g - id\|$$

for some $C_2 > 0$.

The above inequalities give the following

$$|v_1|, |v_2| \leq C'(\delta + \|g - id\|)$$

and

$$v_3 \leq -1 + C''(\delta + \|g - id\|).$$

This gives that

$$|\angle(-j, v)| \leq C'''(\delta + \|g - id\|).$$

The lemma is proved. \square

Let $L = \{(0, 0, t) : t > 0\} \subset \mathbb{H}^3$ be the geodesic through $j = (0, 0, 1) \in \mathbb{H}^3$ with the ideal endpoint $0 \in \mathbf{C} \subset \partial_\infty \mathbb{H}^3$.

Lemma A.3. *Let h be a geodesic in $\mathbb{H}^2 \subset \mathbb{H}^3$ that intersects L between points j and $e^{-r}j$ for some $r \geq 1$. Given $\epsilon_0, \delta_0 > 0$, there exist $C(r, \epsilon_0, \delta_0), C_0(r, \epsilon_0, \delta_0) > 0$ such that*

$$D_{T\mathbb{H}^3}(R_h^\epsilon(\{z, u\}), \{e^{-r}j, -j\}) \leq C_0\delta + C\epsilon$$

for any $0 \leq \epsilon < \epsilon_0$, $0 \leq \delta < \delta_0$, and $\{z, u\} \in T\mathbb{H}^3$ with

$$D_{T\mathbb{H}^3}(\{z, u\}, \{e^{-r}j, -j\}) \leq \delta,$$

where R_h^ϵ is the hyperbolic rotation around the axis h by the angle ϵ .

Proof. The quantity $D_{T\mathbb{H}^3}(R_h^\epsilon(\{z, u\}), \{e^{-r}j, -j\})$ is the largest when h is orthogonal to L at the point j . In this case R_h^ϵ fixes 1 and -1 , and

$$R_h^\epsilon(z) = \frac{\cos \frac{\epsilon}{2} z - i \sin \frac{\epsilon}{2}}{-i \sin \frac{\epsilon}{2} z + \cos \frac{\epsilon}{2}}.$$

Therefore, there exists $C > 0$ such that

$$\|R_h^\epsilon - id\| \leq C\epsilon.$$

The lemma follows by Lemma A.2. \square

The following lemma is standard (cf. [3]).

Lemma A.4. *Let $D_{r_0}(j) \subset \mathbb{H}^3$ be the hyperbolic ball of radius $r_0 > 0$ centered at $j \in \mathbb{H}^3$. Then there exists $C = C(r_0) > 0$ such that if h_1, h_2 are two hyperbolic geodesics with a common endpoint that intersect $D_{r_0}(j)$ and if $d_{r_0}(h_1, h_2)$ is the hyperbolic distance between $h_1 \cap D_{r_0}(j)$ and $h_2 \cap D_{r_0}(j)$ then*

$$\|R_{h_1}^\epsilon R_{h_2}^{-\epsilon} - id\| \leq C d_{r_0}(h_1, h_2) \epsilon$$

for any $\epsilon > 0$.

Lemma A.5. *Let g be a geodesic in $\mathbb{H}^2 \subset \mathbb{H}^3$ that intersects $L = \{(0, 0, t) : t > 0\}$ between j and $e^{-m}j$ at an angle $\epsilon > 0$. Then, for any $\delta > 0$,*

$$D_{T\mathbb{H}^3}(\{R_g^\theta(\{e^{-m}j, -j\}), \{e^{-m}j, -j\}\}) < \delta$$

when $\epsilon = \epsilon(\delta) > 0$ is small enough.

Proof. The angle $\epsilon_1 > 0$ between $R_g^\theta(L)$ and L satisfies

$$(18) \quad |\epsilon_1| \leq \frac{\pi}{2} \cdot |\theta| \cdot |\epsilon|.$$

To see this, we normalize such that $g = L$ and L has turned into a geodesic with endpoints $a < 0$ and $b > 0$. The the geodesic L is parametrized by $\gamma(t) = \frac{a+b}{2} + \frac{b-a}{2} \cos t + (\frac{a+b}{2} + \frac{b-a}{2} \sin t)j$ and $R_g^\theta(L)$ is parametrized by $\gamma_1(t) = (\frac{a+b}{2} + \frac{b-a}{2} \cos t) \cos \theta + (\frac{a+b}{2} + \frac{b-a}{2} \cos t) \sin \theta i + (\frac{a+b}{2} + \frac{b-a}{2} \sin t)j$. Thus

$$\cos \epsilon_1 = \frac{\gamma'(t) \cdot \gamma_1'(t)}{\|\gamma'(t)\| \cdot \|\gamma_1'(t)\|} = \sin^2 \epsilon \cos \theta + \cos^2 \epsilon$$

which implies

$$|\epsilon_1| \leq |2 \sin^{-1}(\epsilon \sin \frac{\theta}{2})| \leq \frac{\pi}{2} |\theta| \cdot |\epsilon|.$$

We go back to the assumption that $L = \{(0, 0, t) : t > 0\}$. By (18), the absolute value of the angle $|\epsilon_1|$ between L and $R_g^\theta(L)$ is less than $\frac{\pi^2}{2} |\epsilon|$. Let $P = g \cap L$. The angle (after the Euclidean parallel transport) between the tangent vector to $R_g^\theta([P, e^{-m'}j])$ at the endpoint $e^{-m'}j$ and the vector $-j$ decreases as $m' \rightarrow m$ for all m' such that $ht(P) \geq ht(e^{-m'}j) \geq ht(e^{-m}j)$. Thus the angle is the largest if $P = e^{-m}j$ which implies that $|\epsilon_1| \leq \frac{\pi^2}{2} |\epsilon|$.

Thus, for R large enough, we have

$$|\angle(R_g^\theta(\{e^{-m}j, -j\}), \{e^{-m}j, -j\})| < \delta.$$

To estimate the height and z -coordinate of $R_g^\theta(e^{-m}j)$, we note that both $|ht(R_h^\theta(e^{-m}j)) - ht(e^{-m}j)|$ and $|Z(R_h^\theta(e^{-m}j))|$ are the largest when $g \cap h = p = j$. Then the angle between g and $R_h^\theta(g)$ is ϵ_1 , where $|\epsilon_1| \leq \frac{\pi^2}{2}|\epsilon|$. Since g and $R_h^\theta(g)$ belong to a hyperbolic plane embedded in \mathbb{H}^3 which contains the g , we can restrict further analysis to the upper half-plane \mathbb{H}^2 where we identify $i \in \mathbb{H}^2$ with $j \in \mathbb{H}^3$. Let $A \in PSL_2(\mathbb{R})$ be an isometry of \mathbb{H}^2 which fixes $i \in \mathbb{H}^2$ and maps g into $R_h^\theta(g)$. Then $A(e^{-m}i) = R_h^\theta(e^{-m}j)$ for the embedding $\mathbb{H}^2 \subset \mathbb{H}^3$. Note that $A(z) = \frac{cz+d}{dz+c}$ with $c, d > 0$ and $c^2 + d^2 = 1$. It follows that $|\frac{d}{c}| \leq C\epsilon_1$, for some $C > 0$ and for $|\epsilon|$ small enough. Furthermore,

$$A(e^{-m}i) = \frac{e^{-m}i + \frac{d}{c}}{-\frac{d}{c}e^{-m}i + 1}$$

which implies that

$$\left| \operatorname{Im}(A(e^{-m}i)) - e^{-m} \right| \leq C_{11} \left(\frac{d}{c} \right)^2 e^{-m}$$

and

$$\left| \operatorname{Re}(A(e^{-m}i)) \right| \leq C_{12} \frac{d}{c}$$

and the lemma follows. \square

REFERENCES

- [1] A. Beardon, *The geometry of discrete groups*, Graduate Texts in Mathematics, 91, Springer-Verlag, New York, 1983.
- [2] F. Bonahon, *Transverse Hölder distributions for geodesic laminations*, Topology 36 (1997), no. 1, 103122.
- [3] F. Bonahon, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, Ann. Fac. Sci. Toulouse Math. (6) 5 (1996), no. 2, 233297.
- [4] L. Bowen, *Weak Forms of the Ehrenpreis Conjecture and the Surface Subgroup Conjecture*, arXiv:math/0411662.
- [5] C. Earle, I. Kra and S. L. Krushkal, *Holomorphic motions and Teichmüller spaces*, Trans. Amer. Math. Soc. 343 (1994), no. 2, 927948.
- [6] D.B.A. Epstein and A. Marden, *Convex hulls in hyperbolic space, a theorem of Sullivan and measured pleated surfaces*, LMS Lecture Notes 111, pages 112-253. Cambridge University Press, 1987.
- [7] D.B.A. Epstein, A. Marden and V. Markovic, *Quasiconformal homeomorphisms and the convex hull boundary*, Ann. of Math. (2) 159 (2004), no. 1, 305336.
- [8] L. Keen and C. Series, *How to bend pairs of punctured tori*, Lipa's legacy (New York, 1995), 359387, Contemp. Math., 211, Amer. Math. Soc., Providence, RI, 1997.
- [9] J. Kahn and V. Markovic, *Immersing almost geodesic surfaces in a closed hyperbolic 3-manifold*, to appear Ann. Math.
- [10] C. Kourouniotis, *Complex length coordinates for quasi-Fuchsian groups*, Mathematica 41 (1994), no. 1, 173-188.
- [11] R. Mañé, P. Sad and D. Sullivan, *On the dynamics of rational maps*, Ann. Sci. Ecole Norm. Sup, 16, 193-217, 1983.
- [12] C. Moore, *Exponential decay of correlation coefficients for geodesic flows*, Group representations, ergodic theory, operators algebras, and mathematical physics, (Berkeley, Calif., 1984), 163-181, Math. Sci. Res. Inst. Publ., 6, Springer, New York, 1987.
- [13] D. Šarić, *Real and Complex Earthquakes*, Trans. Amer. Math. Soc. 358 (2006), no. 1, 233-249.
- [14] D. Šarić, *Bendings by finitely additive transverse cocycles*, available on arXiv.
- [15] Z. Ślodkowski, *Holomorphic motions and polynomial hulls*, Proc. Amer. Math. Soc. 111 (1991), no. 2, 347355.

- [16] S. P. Tan, *Complex Fenchel-Nielsen coordinates for quasi-Fuchsian structures*, Internat. J. Math. 5 (1994), no. 2, 239251.

DEPARTMENT OF MATHEMATICS, QUEENS COLLEGE OF CUNY, 65-30 KISSENA BLVD., FLUSHING, NY 11367

E-mail address: Dragomir.Saric@qc.cuny.edu

MATHEMATICS PH.D. PROGRAM, THE CUNY GRADUATE CENTER, 365 FIFTH AVENUE, NEW YORK, NY 10016-4309