

QUEENS COLLEGE
DEPARTMENT OF MATHEMATICS
FINAL EXAMINATION

MATHEMATICS 142

2½ HOURS

SPRING 2015

INSTRUCTIONS:

ANSWER ALL QUESTIONS

SHOW ALL WORK

1. Evaluate the following integrals without the use of a calculator.

(a) $\int \frac{3z^2}{\sqrt{1+z^3}} dz$

(c) $\int \frac{6+u^2}{2u^3} du$

(b) $\int_0^5 \sqrt{25-x^2} dx$

(d) $\int \frac{3e^{\sqrt{x}}}{\sqrt{x}} dx$

2. Using the definition of a definite integral as the limit of the Riemann sum, evaluate

$$\int_0^2 (2x^2 - 2) dx.$$

Note: $\sum_{k=1}^n k = \frac{n(n+1)}{2}, \quad \sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$

3. If $F(x) = \int_1^x f'(t) dt$ where $f(t) = \int_1^{t^2} (u^3 + 2u^2) du$, find $F'(3)$.

4. In each case, calculate the derivative of the given function.

(a) $f(x) = \frac{(x^3 + 3x)^3 \sin^2 x}{\sqrt[5]{x}}$

(c) $h(x) = x^x$

(b) $g(x) = \cos^{-1}(e^x)$

(d) $k(x) = \frac{\cos^{-1}(3x)}{\ln(3x^2 + 6x)}$

5. Calculate the volume of a cone with height h and base radius r .

(Hint: Rotate an appropriately defined line around the x -axis).

6. Find the volume of the solid obtained by rotating the region bounded by $y = x - x^2$ and $y = 0$ about the line $x = 3$.

7. (a) Show that $f(x) = \frac{4x-1}{2x+3}$ has an inverse $f^{-1}(x)$ and then (b) Calculate the derivative of $f^{-1}(x)$ in two ways:

(i) by differentiating the expression for $f^{-1}(x)$,

(ii) by using the general formula for the derivative of an inverse function.

8. Find the arc length function for the curve $y = f(x) = x^2 - \ln(x)$ taking $P_0(1, 1)$ as the starting point. Find the arc length along the curve from $(1, 1)$ to $(5, f(5))$.

9. Suppose that the growth rate of bacteria in a dish is proportional to the population of bacteria at any time t , where t is in hours (That is, $\frac{dP}{dt} = kP$). Let $P = 25,000$ at $t = 0$ and $P = 42,000$ at $t = 12$. Determine k .

10. Find the solution of the differential equation that satisfies the given initial condition.

$$\frac{dy}{dx} = \frac{e^{2x}(1+y^3)}{y^2}, \quad y(1) = 3$$