

Math 142(2)

Name: _____

Spring 2015

Exam #1

3/9/2015

Time Limit: 75 Minutes

You may *not* use your books or notes on this exam. You are required to show your work on each problem on this exam. The following rules apply:

- **Organize your work**, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- **Mysterious or unsupported answers will not receive full credit.** A correct answer, unsupported by calculations, explanation, or algebraic work will receive no credit; an incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.
- If you need more space, use the back of the pages; clearly indicate when you have done this.

Grade Table (for teacher use only)

Question	Points	Score
1	20	
2	20	
3	20	
4	20	
5	20	
Total:	100	

1. (20 points) Find the most general antiderivative of the following functions. You need not simplify your answer where applicable.

(i)

$$f(x) = (9x^8 + 5x^{2/5}) dx$$

$$F(x) = x^9 + 5 \cdot \frac{5}{7} x^{7/5} + C$$

(ii)

$$g(x) = \frac{2x + x^2 + x^3}{\sqrt{x}} + C$$

$$G(x) = \frac{4}{3} x^{3/2} + \frac{2}{5} x^{5/2} + \frac{2}{7} x^{7/2} + C$$

(iii)

$$h(x) = \cos(x) + 3\sec^2(x)$$

$$H(x) = \sin(x) + 3\tan(x) + C$$

2. (20 points) Using the limit-sum definition for area under a curve, calculate $\int_0^1 (3x^2 + x + 1) dx$.

$$\left(\text{Hint : } \sum_{i=1}^n i = \frac{n(n+1)}{2}, \quad \int_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6} \right)$$

We have:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n \Delta x f(x_i)$$

In our case,

$$\Delta x = \frac{b-a}{n} = \frac{1}{n}, \quad x_i = a + i\Delta x = \frac{i}{n}$$

As such,

$$\begin{aligned} \int_0^1 (3x^2 + x + 2) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left(3 \frac{i^2}{n^2} + \frac{i}{n} + 2 \right) = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{3}{n^2} \sum_{i=1}^n i^2 + \frac{1}{n} \sum_{i=1}^n i + 2 \sum_{i=1}^n 1 \right] \\ &= \lim_{n \rightarrow \infty} \left(\frac{1}{n} \left[\frac{3n(n+1)(2n+1)}{6n^2} + \frac{n(n+1)}{2n} + 2n \right] \right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{3n(n+1)(2n+1)}{6n^3} + \frac{n(n+1)}{2n^2} + 2 \right) \\ &= \frac{6}{6} + \frac{1}{2} + 2 \end{aligned}$$

3. (20 points) (a) (5 points) Let $F'(x) = f(x)$. From the evaluation theorem, complete the following equality:

$$\int_a^b f(x) dx = F(b) - F(a)$$

- (b) (15 points) Evaluate the following integrals:

(i) $\int_{-1}^1 \sqrt{t}(1-t) dt = \left(\frac{2}{3}t^{3/2} - \frac{2}{5}t^{5/2}\right)\Big|_{-1}^1.$

(ii) $\int_0^1 \left(\sqrt{\frac{4}{x}}\right) dx = 4x^{1/2}\Big|_0^1.$

(iii) $\int_{-9}^8 \pi^2 dx = \pi^2(8+9)$

4. (20 points) (a) (5 points) Describe the difference between the following two expressions:

$$\int_2^x f(t) dt \quad \& \quad \int_2^9 f(t) dt$$

The first expression is a function. The second one is a constant, because it's the area under a curve.

- (b) (15 points) Calculate the derivative for $f(x) = \int_2^{x^5} ((t^2 + t) \cos(t)) dt$.

$$\begin{aligned} \frac{d}{dx} \left(\int_2^{x^5} ((t^2 + t) \cos(t)) \right) &\stackrel{u=x^5}{=} \frac{d}{du} \left(\int_2^u ((t^2 + t) \cos(t)) \right) \frac{du}{dx} \\ &= ((u^2 + u) \cos(u)) (5x^4) = ((x^{10} + x^5)(\cos(x^5)))(5x^4) \end{aligned}$$

5. (20 points) Calculate the following antiderivatives.

(i)

$$\int x^2 \sqrt{x^3 + 2} dx$$

Let $u = x^3 + 2$, so then $du = 3x^2$.

$$\int x^2 \sqrt{x^3 + 2} dx = \int \frac{1}{3} \sqrt{u} du = \frac{1}{3} \cdot \frac{2}{3} u^{3/2} = \frac{2}{9} (x^3 + 2)^{3/2}$$

(ii)

$$\int x \sin(x^2 + 2) dx$$

Let $u = x^2 + 2$, $du = 2x$. Then,

$$\int x \sin(x^2 + 2) dx = \int \frac{1}{2} \sin(u) du = \frac{-1}{2} \cos(u) = -\frac{1}{2} \cos(x^2 + 2).$$

(iii)

$$\int \frac{\cos(x)}{2 \sin^2(x)} dx$$

Let $u = \sin(x)$, $du = \cos(x)$.

$$\int \frac{\cos(x)}{2 \sin^2(x)} dx = \int \frac{1}{2u^2} du = -u^{-1} = -\frac{1}{2} (\sin(x))^{-1}$$