

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

*Question:* Does  $\iint_D f(x, y) dy dx$  make sense when domain  $D$  is not a rectangle?

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

*Question:* Does  $\iint_D f(x, y) dy dx$  make sense when domain  $D$  is not a rectangle?

*Answer:* Yes, because we can view  $\iint_D$  as a  $\iint_R$ :

Suppose  $D$  is not a rectangle. Then *fit*  $D$  in a rectangle  $R$ ,

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

*Question:* Does  $\iint_D f(x, y) dy dx$  make sense when domain  $D$  is not a rectangle?

*Answer:* Yes, because we can view  $\iint_D$  as a  $\iint_R$ :

Suppose  $D$  is not a rectangle. Then *fit*  $D$  in a rectangle  $R$ , and extend the function  $f(x, y)$  to be defined over all  $R$ :

$$F(x, y) = \begin{cases} f(x, y) & (x, y) \in D \\ 0 & (x, y) \notin D \end{cases}$$

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

*Question:* Does  $\iint_D f(x, y) dy dx$  make sense when domain  $D$  is not a rectangle?

*Answer:* Yes, because we can view  $\iint_D$  as a  $\iint_R$ :

Suppose  $D$  is not a rectangle. Then *fit*  $D$  in a rectangle  $R$ , and extend the function  $f(x, y)$  to be defined over all  $R$ :

$$F(x, y) = \begin{cases} f(x, y) & (x, y) \in D \\ 0 & (x, y) \notin D \end{cases}$$

Then define  $\iint_D f(x, y) dA = \iint_R F(x, y) dA$ .

## General regions *ARE* rectangles

Last time:  $\iint_R f(x, y) dy dx$  when  $R$  is a rectangle. (Riemann)

*Question:* Does  $\iint_D f(x, y) dy dx$  make sense when domain  $D$  is not a rectangle?

*Answer:* Yes, because we can view  $\iint_D$  as a  $\iint_R$ :

Suppose  $D$  is not a rectangle. Then *fit*  $D$  in a rectangle  $R$ , and extend the function  $f(x, y)$  to be defined over all  $R$ :

$$F(x, y) = \begin{cases} f(x, y) & (x, y) \in D \\ 0 & (x, y) \notin D \end{cases}$$

Then define  $\iint_D f(x, y) dA = \iint_R F(x, y) dA$ .

(Which we know exists)

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,



## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

Type I

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{a "left function" } x = h_2(y) \\ \text{a "right function" } x = h_1(y) \end{array} \right\}$ ,

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{a "left function" } x = h_2(y) \\ \text{a "right function" } x = h_1(y) \end{array} \right\}$ ,

then integrate by slices with fixed  $y$  values.

$$\iint_D f(x, y) dA = \int_{y=c}^{y=d} \int_{x=h_1(y)}^{x=h_2(y)} f(x, y) dx dy$$

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{a "left function" } x = h_2(y) \\ \text{a "right function" } x = h_1(y) \end{array} \right\}$ ,

then integrate by slices with fixed  $y$  values.

$$\iint_D f(x, y) dA = \int_{y=c}^{y=d} \int_{x=h_1(y)}^{x=h_2(y)} f(x, y) dx dy$$

Type I

Type II

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{a "left function" } x = h_2(y) \\ \text{a "right function" } x = h_1(y) \end{array} \right\}$ ,

then integrate by slices with fixed  $y$  values.

$$\iint_D f(x, y) dA = \int_{y=c}^{y=d} \int_{x=h_1(y)}^{x=h_2(y)} f(x, y) dy dx$$

Determine type by looking at which slices cut all the way through  $D$ .

Type I

Type II

## Calculating double integrals over non-rectangles

The way we decide to integrate  $\iint_D$  depends on the shape of  $D$ :

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{an "upper function" } y = g_2(x) \\ \text{a "lower function" } y = g_1(x) \end{array} \right\}$ ,

**Type I**

then integrate by slices with fixed  $x$  values.

$$\iint_D f(x, y) dA = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy dx$$

If  $D$  is defined by  $\left\{ \begin{array}{l} \text{a "left function" } x = h_2(y) \\ \text{a "right function" } x = h_1(y) \end{array} \right\}$ ,

**Type II**

then integrate by slices with fixed  $y$  values.

$$\iint_D f(x, y) dA = \int_{y=c}^{y=d} \int_{x=h_1(y)}^{x=h_2(y)} f(x, y) dx dy$$

Determine type by looking at which slices cut all the way through  $D$ . Some regions work either way. Choose based on  $f(x, y)$ .

## Simple Example

**Example.** Find  $\iint_D (x + 2y) dA$ ,  
where  $D$  is bounded by  $y = 2x^2$  and  $y = 1 + x^2$ .



## Simple Example

**Example.** Find  $\iint_D (x + 2y) dA$ ,  
where  $D$  is bounded by  $y = 2x^2$  and  $y = 1 + x^2$ .

**Steps:**

1. Plot the curves (Draw a picture!)
2. Find points of intersection
3. Determine order of integration
4. Determine “upper” and “lower” functions, other bounds
5. Do the integrals.

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_ and the “lower” function is \_\_\_\_\_.

We calculate  $\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} xy \, dx \, dy =$

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_  
and the “lower” function is \_\_\_\_\_.

We calculate 
$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} xy \, dx \, dy = \int_{y=-2}^{y=4} \left[ y \frac{x^2}{2} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy =$$



## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_ and the “lower” function is \_\_\_\_\_.

$$\text{We calculate } \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} xy \, dx \, dy = \int_{y=-2}^{y=4} \left[ y \frac{x^2}{2} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy =$$

$$\frac{1}{2} \int_{y=-2}^{y=4} y(y+1)^2 - y\left(\frac{y^2-6}{2}\right)^2 dy =$$

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_ and the “lower” function is \_\_\_\_\_.

$$\begin{aligned} \text{We calculate } \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} xy \, dx \, dy &= \int_{y=-2}^{y=4} \left[ y \frac{x^2}{2} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy = \\ \frac{1}{2} \int_{y=-2}^{y=4} y(y+1)^2 - y\left(\frac{y^2-6}{2}\right)^2 dy &= \frac{1}{2} \int_{y=-2}^{y=4} \left(-\frac{y^5}{4} + 4y^3 + 2y^2 - 8y\right) dy \end{aligned}$$

## Not-as-simple Example

**Example.** Find  $\iint_D xy \, dA$ , where  $D$  bdd by  $y = x - 1$  and  $y^2 = 2x + 6$ .

**Important:** Draw a picture.

If we were to set up the integral as slices in  $x$ , there would be two different lower functions, depending on whether  $x \leq 1$  or  $x \geq 1$ .

This would require doing **two** integrals! (What are they?)

Instead, integrate with slices in  $y$ . The “upper” function is \_\_\_\_\_ and the “lower” function is \_\_\_\_\_.

$$\begin{aligned} \text{We calculate } \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} xy \, dx \, dy &= \int_{y=-2}^{y=4} \left[ y \frac{x^2}{2} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy = \\ \frac{1}{2} \int_{y=-2}^{y=4} y(y+1)^2 - y\left(\frac{y^2-6}{2}\right)^2 dy &= \frac{1}{2} \int_{y=-2}^{y=4} \left(-\frac{y^5}{4} + 4y^3 + 2y^2 - 8y\right) dy \\ &= \frac{1}{2} \left[ -\frac{y^6}{24} + y^4 + \frac{2}{3}y^3 - 4y^2 \right]_{y=-2}^{y=4} = 36 \end{aligned}$$

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid. Project the solid onto  $xy$ -plane to find domain  $D$ .

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid.

Project the solid onto  $xy$ -plane to find domain  $D$ .

Where does  $x + 2y + z = 2$  intersect the axes?

(Draw in 3-space and on  $xy$ -plane.)

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid.

Project the solid onto  $xy$ -plane to find domain  $D$ .

Where does  $x + 2y + z = 2$  intersect the axes?

(Draw in 3-space and on  $xy$ -plane.)

What does  $z = 0$  do? What does  $x = 0$  do?

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid.

Project the solid onto  $xy$ -plane to find domain  $D$ .

Where does  $x + 2y + z = 2$  intersect the axes?

(Draw in 3-space and on  $xy$ -plane.)

What does  $z = 0$  do? What does  $x = 0$  do?

What does  $x = 2y$  do?



## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid.

Project the solid onto  $xy$ -plane to find domain  $D$ .

Where does  $x + 2y + z = 2$  intersect the axes?

(Draw in 3-space and on  $xy$ -plane.)

What does  $z = 0$  do? What does  $x = 0$  do?

What does  $x = 2y$  do?

So our domain  $D$  looks like:

(intersection pts? slicing direction? start/stop?)

## A Wordy Example

Sometimes you need to find  $D$  and  $f$  from the problem statement.

**Example.** Set up the integral that finds the volume of the solid bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$ , and  $z = 0$ .

**Solution.** Use the planes to understand and draw the solid.

Project the solid onto  $xy$ -plane to find domain  $D$ .

Where does  $x + 2y + z = 2$  intersect the axes?

(Draw in 3-space and on  $xy$ -plane.)

What does  $z = 0$  do? What does  $x = 0$  do?

What does  $x = 2y$  do?

So our domain  $D$  looks like:

(intersection pts? slicing direction? start/stop?)

Our function is  $f(x, y) = z = 2 - x - 2y$ , and our integral is

$$\int_{x=0}^{x=1} \int_{y=x/2}^{y=1-x/2} (2 - x - 2y) dy dx$$

## Changing the order of integration

We might want to change the order of integration in iterated integrals.

**Caution:** For non-rectangles, we have to be very careful!

## Changing the order of integration

We might want to change the order of integration in iterated integrals.

**Caution:** For non-rectangles, we have to be very careful!

$$\int_0^1 \int_{y=x}^{y=1} f(x, y) dy dx \rightsquigarrow$$

## Changing the order of integration

We might want to change the order of integration in iterated integrals.


**Caution:** For non-rectangles, we have to be very careful!

$$\int_0^1 \int_{y=x}^{y=1} f(x, y) dy dx \rightsquigarrow$$

## Changing the order of integration

We might want to change the order of integration in iterated integrals.

**Caution:** For non-rectangles, we have to be very careful!




When chopping in  $x$ ,

$$\left\{ \begin{array}{l} x \text{ varies from } 0 \text{ to } 1, \\ \text{upper fcn is } y = 1 \\ \text{lower fcn is } y = x \end{array} \right\}$$

## Changing the order of integration

We might want to change the order of integration in iterated integrals.

**Caution:** For non-rectangles, we have to be very careful!



When chopping in  $x$ ,

$$\left\{ \begin{array}{l} x \text{ varies from } 0 \text{ to } 1, \\ \text{upper fcn is } y = 1 \\ \text{lower fcn is } y = x \end{array} \right\}$$

$\longrightarrow$


When chopping in  $y$ ,

$$\left\{ \begin{array}{l} y \text{ varies from } 0 \text{ to } 1, \\ \text{upper fcn is } x = y \\ \text{lower fcn is } x = 0 \end{array} \right\}$$

## Changing the order of integration

We might want to change the order of integration in iterated integrals.

**Caution:** For non-rectangles, we have to be very careful!

$$\int_0^1 \int_{y=x}^{y=1} f(x, y) dy dx \rightsquigarrow \int_0^1 \int_{x=0}^{x=y} f(x, y) dx dy$$


When chopping in  $x$ ,

$$\left\{ \begin{array}{l} x \text{ varies from } 0 \text{ to } 1, \\ \text{upper fcn is } y = 1 \\ \text{lower fcn is } y = x \end{array} \right\}$$

$\longrightarrow$

When chopping in  $y$ ,

$$\left\{ \begin{array}{l} y \text{ varies from } 0 \text{ to } 1, \\ \text{upper fcn is } x = y \\ \text{lower fcn is } x = 0 \end{array} \right\}$$



## Double integral properties

**Property.** Suppose that  $D = D_1 \cup D_2$  (where  $D_1$  and  $D_2$  don't overlap).

Then

$$\iint_D f \, dA = \iint_{D_1} f \, dA + \iint_{D_2} f \, dA.$$

## Double integral properties

**Property.** Suppose that  $D = D_1 \cup D_2$  (where  $D_1$  and  $D_2$  don't overlap).

Then

$$\iint_D f \, dA = \iint_{D_1} f \, dA + \iint_{D_2} f \, dA.$$

**Consequence:** Break down complicated regions into Type I and Type II regions.

## Double integral properties

**Property.** Suppose that  $D = D_1 \cup D_2$  (where  $D_1$  and  $D_2$  don't overlap). Then

$$\iint_D f \, dA = \iint_{D_1} f \, dA + \iint_{D_2} f \, dA.$$

**Consequence:** Break down complicated regions into Type I and Type II regions.

**Property.** Suppose  $m \leq f(x, y) \leq M$  for all  $(x, y) \in D$ . Then

$$m \cdot \text{Area}(D) \leq \iint_D f(x, y) \, dA \leq M \cdot \text{Area}(D)$$

## Double integral properties

**Property.** Suppose that  $D = D_1 \cup D_2$  (where  $D_1$  and  $D_2$  don't overlap). Then

$$\iint_D f \, dA = \iint_{D_1} f \, dA + \iint_{D_2} f \, dA.$$

**Consequence:** Break down complicated regions into Type I and Type II regions.

**Property.** Suppose  $m \leq f(x, y) \leq M$  for all  $(x, y) \in D$ . Then

$$m \cdot \text{Area}(D) \leq \iint_D f(x, y) \, dA \leq M \cdot \text{Area}(D)$$

**Consequence:** This gives a crude approximation for the integral.

## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.


## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

---

mass density function  $\rho(x, y)$


(mass per  
unit area)



## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

mass density function  $\rho(x, y)$   
(mass per  
unit area)




The total mass of the object is

$$m = \iint_D \rho(x, y) dA$$

## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.


mass density function  $\rho(x, y)$   
(mass per  
unit area)



The total mass of the object is

$$m = \iint_D \rho(x, y) dA$$



charge density function  $\sigma(x, y)$   
(charge per  
unit area)







## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

mass density function $\rho(x, y)$ (mass per unit area) 	The total mass of the object is $m = \iint_D \rho(x, y) dA$
charge density function $\sigma(x, y)$ (charge per unit area) 	The total charge on the object is $Q = \iint_D \sigma(x, y) dA$

## Application: Density



Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

mass density function $\rho(x, y)$ (mass per unit area) 	The total mass of the object is $m = \iint_D \rho(x, y) dA$
charge density function $\sigma(x, y)$ (charge per unit area) 	The total charge on the object is $Q = \iint_D \sigma(x, y) dA$

**Example.** Find the mass of a  $\triangle$  lamina w/ corners  $(1, 0)$ ,  $(0, 2)$ ,  $(1, 2)$ , and mass density function  $\rho(x, y) = 1 + 3x + y$ .

## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.



mass density function $\rho(x, y)$ (mass per unit area) 	The total mass of the object is $m = \iint_D \rho(x, y) dA$
charge density function $\sigma(x, y)$ (charge per unit area) 	The total charge on the object is $Q = \iint_D \sigma(x, y) dA$

**Example.** Find the mass of a  $\triangle$  lamina w/ corners  $(1, 0)$ ,  $(0, 2)$ ,  $(1, 2)$ , and mass density function  $\rho(x, y) = 1 + 3x + y$ .

**Solution.**  $m = \int_{x=0}^{x=1} \int_{y=2-2x}^{y=2} (1 + 3x + y) dy dx$

## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

mass density function $\rho(x, y)$ (mass per unit area) 	The total mass of the object is $m = \iint_D \rho(x, y) dA$
charge density function $\sigma(x, y)$ (charge per unit area) 	The total charge on the object is $Q = \iint_D \sigma(x, y) dA$

**Example.** Find the mass of a  $\triangle$  lamina w/ corners  $(1, 0)$ ,  $(0, 2)$ ,  $(1, 2)$ , and mass density function  $\rho(x, y) = 1 + 3x + y$ .



**Solution.**

$$m = \int_{x=0}^{x=1} \int_{y=2-2x}^{y=2} (1 + 3x + y) dy dx$$

$$= \int_{x=0}^{x=1} \left( y + 3xy + \frac{y^2}{2} \right) \Big|_{y=2-2x}^{y=2} dx$$

## Application: Density

Suppose you have a 2D sheet of metal (a **lamina**) where density varies over the sheet.

mass density function $\rho(x, y)$ (mass per unit area) 	The total mass of the object is $m = \iint_D \rho(x, y) dA$
charge density function $\sigma(x, y)$ (charge per unit area) 	The total charge on the object is $Q = \iint_D \sigma(x, y) dA$

**Example.** Find the mass of a  $\triangle$  lamina w/ corners  $(1, 0)$ ,  $(0, 2)$ ,  $(1, 2)$ , and mass density function  $\rho(x, y) = 1 + 3x + y$ .

**Solution.**

$$\begin{aligned}
 m &= \int_{x=0}^{x=1} \int_{y=2-2x}^{y=2} (1 + 3x + y) dy dx \\
 &= \int_{x=0}^{x=1} \left( y + 3xy + \frac{y^2}{2} \right) \Big|_{y=2-2x}^{y=2} dx \\
 &= \int_{x=0}^{x=1} (6x + 4x^2) dx = 3x^2 + \frac{4}{3}x^3 \Big|_{x=0}^{x=1} = \frac{13}{3}
 \end{aligned}$$