

Vectors

We will be using vectors and matrices to store and manipulate data.

Definition: A **vector** \vec{v} is a column of numbers. Use bold faced letters or vector signs to distinguish vectors from other variables.

We refer to the **entries** of a vector by using subscripts.

The **length** of a vector is the number of entries it has. (normally n)

Example. $\vec{v} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = [1 \ 2 \ 3]^T.$

Vectors

We will be using vectors and matrices to store and manipulate data.

Definition: A **vector** \vec{v} is a column of numbers. Use bold faced letters or vector signs to distinguish vectors from other variables.

We refer to the **entries** of a vector by using subscripts.

The **length** of a vector is the number of entries it has. (normally n)

Example. $\vec{v} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = [1 \ 2 \ 3]^T.$

Example. Use a vector to represent the age distribution of a population: let F_i be the number of females with ages in the interval $[5i, 5(i+1))$. We can represent the total female population by the vector \vec{F} .

The females from 0 up to 5 are counted in F_0 ;
those from 5 up to 10 are counted in F_1 , etc.

$$\vec{F} = \begin{bmatrix} F_0 \\ F_1 \\ F_2 \\ \vdots \\ F_{n-1} \end{bmatrix}$$

Matrices

Definition: A **matrix** A is a two-dimensional array of numbers.

A matrix with m rows and n columns is called an $m \times n$ matrix.

★ Row by column — Row by column — Row by column ★

Note: A vector can be thought of as an $n \times 1$ matrix.

Matrices

Definition: A **matrix** A is a two-dimensional array of numbers.

A matrix with m rows and n columns is called an $m \times n$ matrix.

★ Row by column — Row by column — Row by column ★

Note: A vector can be thought of as an $n \times 1$ matrix.

Matrices are denoted by a capital letter. Entries are lower case and have two subscripts, the corresponding row and column.

Example. A generic 2×3 matrix has the form $A = \begin{bmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \end{bmatrix}$.

Matrices

Definition: A **matrix** A is a two-dimensional array of numbers.

A matrix with m rows and n columns is called an $m \times n$ matrix.

★ Row by column — Row by column — Row by column ★

Note: A vector can be thought of as an $n \times 1$ matrix.

Matrices are denoted by a capital letter. Entries are lower case and have two subscripts, the corresponding row and column.

Example. A generic 2×3 matrix has the form $A = \begin{bmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \end{bmatrix}$.

Definition: The matrix $B = \begin{bmatrix} 30 & 50 \\ 100 & 250 \end{bmatrix}$ is a **square matrix** because it has the same number of rows as columns.

Matrices

Example. We will sometimes interpret a matrix as a **transition** matrix. In this case, the matrix is square (say $n \times n$), where the n rows and n columns correspond to certain **states** (situations). An entry $a_{i,j}$ represents transitioning from state j to state i .

Matrices

Example. We will sometimes interpret a matrix as a **transition** matrix. In this case, the matrix is square (say $n \times n$), where the n rows and n columns correspond to certain **states** (situations).

An entry $a_{i,j}$ represents transitioning from state j to state i .

Example. In our population example, suppose we want to model people getting older, transitioning from one state (age group) to the next. We would set up a transition matrix such as:

	FROM state:	
TO state:	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$	because everyone in the first age group will move to the second age group ($a_{2,1}$), everyone in state 2 will move to state 3 ($a_{3,2}$), etc.

Matrix Multiplication

The power of matrices arises in their multiplication.

Given two matrices, A of size $m \times k$ and B of size $l \times n$, we can find the product AB **if and only if** k equals l .

Let A be an $m \times k$ matrix and B , $k \times n$. Then AB is of size $m \times n$.

Matrix Multiplication

The power of matrices arises in their multiplication.

Given two matrices, A of size $m \times k$ and B of size $l \times n$, we can find the product AB **if and only if** k equals l .

Let A be an $m \times k$ matrix and B , $k \times n$. Then AB is of size $m \times n$.

To calculate the entries of AB , remember: “Row by column”:

$$\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 0 & 6 \\ -4 & 1 & 2 \end{bmatrix} = \begin{bmatrix} \textcircled{0} & \textcircled{0} & \textcircled{0} \\ \textcircled{0} & \textcircled{0} & \textcircled{0} \end{bmatrix}$$

Matrix Multiplication

The power of matrices arises in their multiplication.

Given two matrices, A of size $m \times k$ and B of size $l \times n$, we can find the product AB **if and only if** k equals l .

Let A be an $m \times k$ matrix and B , $k \times n$. Then AB is of size $m \times n$.

To calculate the entries of AB , remember: "Row by column":

$$\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 0 & 6 \\ -4 & 1 & 2 \end{bmatrix} = \begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$$

When we write A^2 , this means AA ; A^3 means AAA , etc.

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}^2 = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & \circ & \circ \\ 0 & 1 & \circ \\ 0 & 0 & 1 \end{bmatrix}$$

The power of transition matrices

Example. Modeling a changing population using a matrix model.

Let us choose a size of age interval $\Delta=5$ years (“Delta”), and divide the female population into states:

State 0: ages $[0, 5)$ with $F_0 = 150$ females

State 1: ages $[5, 10)$ with $F_1 = 200$ females

State 2: ages $[10, 15)$ with $F_2 = 180$ females

State 3: ages $[15, 20)$ with $F_3 = 120$ females

State 4: ages $[20, 25)$ with $F_4 = 60$ females

The power of transition matrices

Example. Modeling a changing population using a matrix model.

Let us choose a size of age interval $\Delta=5$ years (“Delta”), and divide the female population into states:

State 0: ages $[0, 5)$ with $F_0 = 150$ females

State 1: ages $[5, 10)$ with $F_1 = 200$ females

State 2: ages $[10, 15)$ with $F_2 = 180$ females

State 3: ages $[15, 20)$ with $F_3 = 120$ females

State 4: ages $[20, 25)$ with $F_4 = 60$ females

age distribution vector:

$$\vec{F} = \begin{bmatrix} 150 \\ 200 \\ 180 \\ 120 \\ 60 \end{bmatrix}$$

The power of transition matrices

Example. Modeling a changing population using a matrix model.

Let us choose a size of age interval $\Delta=5$ years (“Delta”), and divide the female population into states:

State 0: ages $[0, 5)$ with $F_0 = 150$ females

State 1: ages $[5, 10)$ with $F_1 = 200$ females

State 2: ages $[10, 15)$ with $F_2 = 180$ females

State 3: ages $[15, 20)$ with $F_3 = 120$ females

State 4: ages $[20, 25)$ with $F_4 = 60$ females

age distribution vector:

$$\vec{F} = \begin{bmatrix} 150 \\ 200 \\ 180 \\ 120 \\ 60 \end{bmatrix}$$

Using a transition matrix, we can determine the population in 5 years:

$$A \cdot \vec{F} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}^1 \begin{bmatrix} 150 \\ 200 \\ 180 \\ 120 \\ 60 \end{bmatrix} = \begin{bmatrix} 0 \\ 150 \\ 200 \\ 180 \\ 120 \end{bmatrix}$$

Leslie Matrices

The transition matrix in the previous example is not entirely realistic, because people die and are born

To take death into account, modify:

Leslie Matrices

The transition matrix in the previous example is not entirely realistic, because people die and are born

To take death into account, modify:

To take birth into account, modify:

(j females !)

Leslie Matrices

The transition matrix in the previous example is not entirely realistic, because people die and are born

To take death into account, modify:

To take birth into account, modify:

(j females !)

The resulting transition matrix is called a **Leslie matrix**:

Let m_i be the average number of females that women in state i bear.

Let p_i be the fraction of women in state i that survive to state $i + 1$.

Leslie Matrices

The transition matrix in the previous example is not entirely realistic, because people die and are born

To take death into account, modify:

To take birth into account, modify:

(j females !)

The resulting transition matrix is called a **Leslie matrix**:

Let m_i be the average number of females that women in state i bear.

Let p_i be the fraction of women in state i that survive to state $i + 1$.

$$\text{then } \begin{bmatrix} F_0(t + \Delta) \\ F_1(t + \Delta) \\ F_2(t + \Delta) \\ \vdots \\ F_{n-1}(t + \Delta) \end{bmatrix} = \begin{bmatrix} m_0 & m_1 & m_2 & \cdots & m_{n-1} \\ p_0 & 0 & 0 & \cdots & 0 \\ 0 & p_1 & 0 & \cdots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & \cdots & 0 & p_{n-2} & 0 \end{bmatrix} \begin{bmatrix} F_0(t) \\ F_1(t) \\ F_2(t) \\ \vdots \\ F_{n-1}(t) \end{bmatrix}$$

$$\vec{F}(t + \Delta) = M \cdot \vec{F}(t)$$

Leslie Matrices

Example. An animal population example (p. 47)

The population in three age groups, $F_0 = 80$, $F_1 = 40$, and $F_2 = 20$.

Suppose that as Δ time passes, everyone in state 2 dies, and one quarter of everyone else dies. Also suppose that the age-specific maternity rates are $m_0 = 0$, $m_1 = 1$, and $m_2 = 2$. Determine the Leslie matrix and the population distributions at times Δ and 2Δ .

$$\begin{bmatrix} & & \\ & 0 & 0 \\ 0 & & 0 \end{bmatrix} \begin{bmatrix} 80 \\ 40 \\ 20 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix} = \vec{F}(\Delta)$$

$$\begin{bmatrix} & & \\ & 0 & 0 \\ 0 & & 0 \end{bmatrix} \begin{bmatrix} \\ \\ \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix} = \vec{F}(2\Delta)$$

Matrices and Vectors in Mathematica

We can use Mathematica to do these calculations. (See Tutorial 5.)

- ▶ Matrices are defined as lists of row vectors.

```
matrix = {{0, 1, 2}, {.75, 0, 0}, {0, .75, 0}}
```

```
vector = {{80}, {40}, {20}}
```

Matrices and Vectors in Mathematica

We can use Mathematica to do these calculations. (See Tutorial 5.)

- ▶ Matrices are defined as lists of row vectors.

```
matrix = {{0, 1, 2}, {.75, 0, 0}, {0, .75, 0}}  
vector = {{80}, {40}, {20}}
```

- ▶ Multiply matrices by using a period (.)

```
matrix.vector
```

- ▶ Find powers of matrices using `MatrixPower`, not `^`
`MatrixPower[matrix, 2]`

Matrices and Vectors in Mathematica

We can use Mathematica to do these calculations. (See Tutorial 5.)

- ▶ Matrices are defined as lists of row vectors.

```
matrix = {{0, 1, 2}, {.75, 0, 0}, {0, .75, 0}}  
vector = {{80}, {40}, {20}}
```

- ▶ Multiply matrices by using a period (.)

```
matrix.vector
```

- ▶ Find powers of matrices using `MatrixPower`, not `^`

```
MatrixPower[matrix, 2]
```

- ▶ So, to find animal population over time, use the code:

```
Table[  
    , {i, 0, 10}]
```