

Introduction to Bijections

Key tool: A useful method of proving that two sets A and B are of the same size is by way of a *bijection*.

A **bijection** is a function or rule that pairs up elements of A and B .

Introduction to Bijections

Key tool: A useful method of proving that two sets A and B are of the same size is by way of a *bijection*.

A **bijection** is a function or rule that pairs up elements of A and B .

Example. The set A of subsets of $\{s_1, s_2, s_3\}$ are in bijection with the set B of binary words of length 3.

Set A: $\{ \emptyset, \{s_1\}, \{s_2\}, \{s_1, s_2\}, \{s_3\}, \{s_1, s_3\}, \{s_2, s_3\}, \{s_1, s_2, s_3\} \}$

Set B: $\{ 000, 100, 010, 110, 001, 101, 011, 111 \}$

Introduction to Bijections

Key tool: A useful method of proving that two sets A and B are of the same size is by way of a *bijection*.

A **bijection** is a function or rule that pairs up elements of A and B .

Example. The set A of subsets of $\{s_1, s_2, s_3\}$ are in bijection with the set B of binary words of length 3.

$$\begin{array}{l}
 \text{Set A: } \{ \emptyset, \{s_1\}, \{s_2\}, \{s_1, s_2\}, \{s_3\}, \{s_1, s_3\}, \{s_2, s_3\}, \{s_1, s_2, s_3\} \} \\
 \text{Bijection: } \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \\
 \text{Set B: } \{ 000, 100, 010, 110, 001, 101, 011, 111 \}
 \end{array}$$

Introduction to Bijections

Key tool: A useful method of proving that two sets A and B are of the same size is by way of a *bijection*.

A **bijection** is a function or rule that pairs up elements of A and B .

Example. The set A of subsets of $\{s_1, s_2, s_3\}$ are in bijection with the set B of binary words of length 3.

$$\begin{array}{l}
 \text{Set A: } \{ \emptyset, \{s_1\}, \{s_2\}, \{s_1, s_2\}, \{s_3\}, \{s_1, s_3\}, \{s_2, s_3\}, \{s_1, s_2, s_3\} \} \\
 \text{Bijection: } \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \quad \updownarrow \\
 \text{Set B: } \{ 000, 100, 010, 110, 001, 101, 011, 111 \}
 \end{array}$$


Rule: Given $a \in A$, (a is a subset), define $b \in B$ (b is a word):
 If $s_i \in a$, then letter i in b is 1. If $s_i \notin a$, then letter i in b is 0.

Introduction to Bijections

Key tool: A useful method of proving that two sets A and B are of the same size is by way of a *bijection*.

A **bijection** is a function or rule that pairs up elements of A and B .

Example. The set A of subsets of $\{s_1, s_2, s_3\}$ are in bijection with the set B of binary words of length 3.



Rule: Given $a \in A$, (a is a subset), define $b \in B$ (b is a word):
 If $s_i \in a$, then letter i in b is 1. If $s_i \notin a$, then letter i in b is 0.

Difficulties:

- ▶ **Finding** the function or rule (requires rearranging, ordering)
- ▶ **Proving** the function or rule (show it **IS** a bijection).

What is a Function?

Reminder: A **function** f from A to B (write $f : A \rightarrow B$) is a rule where for each element $a \in A$, $f(a)$ is defined as an element $b \in B$ (write $f : a \mapsto b$).

What is a Function?

Reminder: A **function** f from A to B (write $f : A \rightarrow B$) is a rule where for each element $a \in A$, $f(a)$ is defined as an element $b \in B$ (write $f : a \mapsto b$).

- ▶ A is called the **domain**. (We write $A = \text{dom}(f)$)
- ▶ B is called the **codomain**. (We write $B = \text{cod}(f)$)

What is a Function?

Reminder: A **function** f from A to B (write $f : A \rightarrow B$) is a rule where for each element $a \in A$, $f(a)$ is defined as an element $b \in B$ (write $f : a \mapsto b$).

- ▶ A is called the **domain**. (We write $A = \text{dom}(f)$)
- ▶ B is called the **codomain**. (We write $B = \text{cod}(f)$)
- ▶ The **range** of f is the set of values that f takes on:

$$\text{rng}(f) = \{b \in B : f(a) = b \text{ for at least one } a \in A\}$$

What is a Function?

Reminder: A **function** f from A to B (write $f : A \rightarrow B$) is a rule where for each element $a \in A$, $f(a)$ is defined as an element $b \in B$ (write $f : a \mapsto b$).

- ▶ A is called the **domain**. (We write $A = \text{dom}(f)$)
- ▶ B is called the **codomain**. (We write $B = \text{cod}(f)$)
- ▶ The **range** of f is the set of values that f takes on:

$$\text{rng}(f) = \{b \in B : f(a) = b \text{ for at least one } a \in A\}$$

Example. Let A be the set of 3-subsets of $[n]$ and let B be the set of 3-lists of $[n]$. Then define $f : A \rightarrow B$ to be the function that takes a 3-subset $\{i_1, i_2, i_3\} \in A$ (with $i_1 \leq i_2 \leq i_3$) to the word $i_1 i_2 i_3 \in B$.

Question: Is $\text{rng}(f) = B$?

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when
For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

“When the inputs are different, the outputs are different.” (picture)

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

“When the inputs are different, the outputs are different.” (picture)

Definition: A function $f : A \rightarrow B$ is **onto** (a **surjection**) when

For each $b \in B$, there exists some $a \in A$ such that $f(a) = b$.

“Every output gets hit.”

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

“When the inputs are different, the outputs are different.” (picture)

Definition: A function $f : A \rightarrow B$ is **onto** (a **surjection**) when

For each $b \in B$, there exists some $a \in A$ such that $f(a) = b$.

“Every output gets hit.”

Definition: A function $f : A \rightarrow B$ is a **bijection** if it is both one-to-one and onto.

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

“When the inputs are different, the outputs are different.” (picture)

Definition: A function $f : A \rightarrow B$ is **onto** (a **surjection**) when

For each $b \in B$, there exists some $a \in A$ such that $f(a) = b$.

“Every output gets hit.”

Definition: A function $f : A \rightarrow B$ is a **bijection** if it is both one-to-one and onto.

The function from the previous page is _____.

What is a Bijection?

Definition: A function $f : A \rightarrow B$ is **one-to-one** (an **injection**) when

For each $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$, then $a_1 = a_2$.

Equivalently,

For each $a_1, a_2 \in A$, if $a_1 \neq a_2$, then $f(a_1) \neq f(a_2)$.

“When the inputs are different, the outputs are different.” (picture)

Definition: A function $f : A \rightarrow B$ is **onto** (a **surjection**) when

For each $b \in B$, there exists some $a \in A$ such that $f(a) = b$.

“Every output gets hit.”

Definition: A function $f : A \rightarrow B$ is a **bijection** if it is both one-to-one and onto.

The function from the previous page is _____.

What is an example of a function that is onto and not one-to-one?

Proving a Bijection

Example. Use a bijection to prove that $\binom{n}{k} = \binom{n}{n-k}$ for $0 \leq k \leq n$.

Proving a Bijection

Example. Use a bijection to prove that $\binom{n}{k} = \binom{n}{n-k}$ for $0 \leq k \leq n$.

Proof. Let A be the set of k -subsets of $[n]$
and let B be the set of $(n - k)$ -subsets of $[n]$.

A bijection between A and B will prove $\binom{n}{k} = |A| = |B| = \binom{n}{n-k}$.

Proving a Bijection

Example. Use a bijection to prove that $\binom{n}{k} = \binom{n}{n-k}$ for $0 \leq k \leq n$.

Proof. Let A be the set of k -subsets of $[n]$
and let B be the set of $(n - k)$ -subsets of $[n]$.

A bijection between A and B will prove $\binom{n}{k} = |A| = |B| = \binom{n}{n-k}$.

Step 1: Find a candidate bijection.

Strategy. Try out a small (enough) example. Try $n = 5$ and $k = 2$.

$$\left\{ \begin{array}{l} \{1, 2\}, \{1, 3\} \\ \{1, 4\}, \{1, 5\} \\ \{2, 3\}, \{2, 4\} \\ \{2, 5\}, \{3, 4\} \\ \{3, 5\}, \{4, 5\} \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \{1, 2, 3\}, \{1, 2, 4\} \\ \{1, 2, 5\}, \{1, 3, 4\} \\ \{1, 3, 5\}, \{1, 4, 5\} \\ \{2, 3, 4\}, \{2, 3, 5\} \\ \{2, 4, 5\}, \{3, 4, 5\} \end{array} \right\}$$

Proving a Bijection

Example. Use a bijection to prove that $\binom{n}{k} = \binom{n}{n-k}$ for $0 \leq k \leq n$.

Proof. Let A be the set of k -subsets of $[n]$
and let B be the set of $(n - k)$ -subsets of $[n]$.

A bijection between A and B will prove $\binom{n}{k} = |A| = |B| = \binom{n}{n-k}$.

Step 1: Find a candidate bijection.

Strategy. Try out a small (enough) example. Try $n = 5$ and $k = 2$.

$$\left\{ \begin{array}{l} \{1, 2\}, \{1, 3\} \\ \{1, 4\}, \{1, 5\} \\ \{2, 3\}, \{2, 4\} \\ \{2, 5\}, \{3, 4\} \\ \{3, 5\}, \{4, 5\} \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \{1, 2, 3\}, \{1, 2, 4\} \\ \{1, 2, 5\}, \{1, 3, 4\} \\ \{1, 3, 5\}, \{1, 4, 5\} \\ \{2, 3, 4\}, \{2, 3, 5\} \\ \{2, 4, 5\}, \{3, 4, 5\} \end{array} \right\}$$

Guess: Let S be a k -subset of $[n]$. Perhaps $f(S) = \underline{\hspace{2cm}}$.

Proving a Bijection

Step 2: Prove f is well defined.

The function f is well defined. If S is any k -subset of $[n]$, then S^c is a subset of $[n]$ with $n - k$ members. Therefore $f : A \rightarrow B$.

Proving a Bijection

Step 2: Prove f is well defined.

The function f is well defined. If S is any k -subset of $[n]$, then S^c is a subset of $[n]$ with $n - k$ members. Therefore $f : A \rightarrow B$.

Step 3: Prove f is a bijection.

Strategy. Prove that f is both one-to-one and onto.

Proving a Bijection

Step 2: Prove f is well defined.

The function f is well defined. If S is any k -subset of $[n]$, then S^c is a subset of $[n]$ with $n - k$ members. Therefore $f : A \rightarrow B$.

Step 3: Prove f is a bijection.

Strategy. Prove that f is both one-to-one and onto.

f is 1-to-1: Suppose that S_1 and S_2 are two k -subsets of $[n]$ such that $f(S_1) = f(S_2)$. That is, $S_1^c = S_2^c$. This means that for all $i \in [n]$, then $i \notin S_1$ if and only if $i \notin S_2$. Therefore $S_1 = S_2$ and f is 1-to-1.

Proving a Bijection

Step 2: Prove f is well defined.

The function f is well defined. If S is any k -subset of $[n]$, then S^c is a subset of $[n]$ with $n - k$ members. Therefore $f : A \rightarrow B$.

Step 3: Prove f is a bijection.

Strategy. Prove that f is both one-to-one and onto.

f is 1-to-1: Suppose that S_1 and S_2 are two k -subsets of $[n]$ such that $f(S_1) = f(S_2)$. That is, $S_1^c = S_2^c$. This means that for all $i \in [n]$, then $i \notin S_1$ if and only if $i \notin S_2$. Therefore $S_1 = S_2$ and f is 1-to-1.

f is onto: Suppose that $T \in B$ is an $(n - k)$ -subset of $[n]$.

We must find a set $S \in A$ satisfying $f(S) = T$. Choose $S = \underline{\hspace{2cm}}$. Then $S \in A$ (why?), and $f(S) = S^c = T$, so f is onto.

We conclude that f is a bijection and therefore, $\binom{n}{k} = \binom{n}{n-k}$.

Alternative methods to prove bijections

Prove that a rule f is a bijection by finding f 's **inverse**:

- ▶ Determine a rule for a candidate inverse function g .

Alternative methods to prove bijections

Prove that a rule f is a bijection by finding f 's **inverse**:

- ▶ Determine a rule for a candidate inverse function g .
- ▶ Show that f is a well defined function **from** A to B .
- ▶ Show that g is a well defined function **from** B to A .

Alternative methods to prove bijections

Prove that a rule f is a bijection by finding f 's **inverse**:

- ▶ Determine a rule for a candidate inverse function g .
- ▶ Show that f is a well defined function **from** A to B .
- ▶ Show that g is a well defined function **from** B to A .
- ▶ Show for all $a \in A$, $g(f(a)) = a$
and for all $b \in B$, $f(g(b)) = b$

Alternative methods to prove bijections

Prove that a rule f is a bijection by finding f 's **inverse**:

- ▶ Determine a rule for a candidate inverse function g .
- ▶ Show that f is a well defined function **from** A to B .
- ▶ Show that g is a well defined function **from** B to A .
- ▶ Show for all $a \in A$, $g(f(a)) = a$
and for all $b \in B$, $f(g(b)) = b$

When g is the inverse of f , both f and g are bijections.

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

$$\begin{aligned} \text{even: } & \{ \emptyset, \{s_1, s_2\}, \{s_1, s_3\}, \{s_2, s_3\} \} \\ \text{odd: } & \{ \{s_1\}, \{s_2\}, \{s_3\}, \{s_1, s_2, s_3\} \} \end{aligned}$$

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

$$\begin{aligned} \text{even: } & \{ \emptyset, \{s_1, s_2\}, \{s_1, s_3\}, \{s_2, s_3\} \} \\ \text{odd: } & \{ \{s_1\}, \{s_2\}, \{s_3\}, \{s_1, s_2, s_3\} \} \end{aligned}$$

Proof. Let A be the set of even-sized subsets of $[n]$ and let B be the set of odd-sized subsets of $[n]$. Consider the function

$$f(S) = \begin{cases} S - \{1\} & \text{if } 1 \in S \\ S \cup \{1\} & \text{if } 1 \notin S \end{cases}.$$

► f is a well defined function from A to B (why?).

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

$$\begin{aligned} \text{even: } & \{ \emptyset, \{s_1, s_2\}, \{s_1, s_3\}, \{s_2, s_3\} \} \\ \text{odd: } & \{ \{s_1\}, \{s_2\}, \{s_3\}, \{s_1, s_2, s_3\} \} \end{aligned}$$

Proof. Let A be the set of even-sized subsets of $[n]$ and let B be the set of odd-sized subsets of $[n]$. Consider the function

$$f(S) = \begin{cases} S - \{1\} & \text{if } 1 \in S \\ S \cup \{1\} & \text{if } 1 \notin S \end{cases}.$$

- ▶ f is a well defined function from A to B (why?).
- ▶ f is also a well defined function from B to A (why?).

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

$$\begin{aligned} \text{even: } & \{ \emptyset, \{s_1, s_2\}, \{s_1, s_3\}, \{s_2, s_3\} \} \\ \text{odd: } & \{ \{s_1\}, \{s_2\}, \{s_3\}, \{s_1, s_2, s_3\} \} \end{aligned}$$

Proof. Let A be the set of even-sized subsets of $[n]$ and let B be the set of odd-sized subsets of $[n]$. Consider the function

$$f(S) = \begin{cases} S - \{1\} & \text{if } 1 \in S \\ S \cup \{1\} & \text{if } 1 \notin S \end{cases}.$$

- ▶ f is a well defined function from A to B (why?).
- ▶ f is also a well defined function from B to A (why?).
- ▶ f^2 is the identity function.

Therefore, f is a bijection, proving the statement, as desired.

Using the inverse function

Example. There exists as many even-sized subsets of $[n]$ as odd-sized subsets of $[n]$.

$$\begin{aligned} \text{even: } & \{ \emptyset, \{s_1, s_2\}, \{s_1, s_3\}, \{s_2, s_3\} \} \\ \text{odd: } & \{ \{s_1\}, \{s_2\}, \{s_3\}, \{s_1, s_2, s_3\} \} \end{aligned}$$

Proof. Let A be the set of even-sized subsets of $[n]$ and let B be the set of odd-sized subsets of $[n]$. Consider the function

$$f(S) = \begin{cases} S - \{1\} & \text{if } 1 \in S \\ S \cup \{1\} & \text{if } 1 \notin S \end{cases}.$$

- ▶ f is a well defined function from A to B (why?).
- ▶ f is also a well defined function from B to A (why?).
- ▶ f^2 is the identity function.

Therefore, f is a bijection, proving the statement, as desired.

Eyebrow-Raising Consequence: $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$