


# Principle of Inclusion-Exclusion

**Example.** Suppose that in this class, 14 students play soccer and 11 students play basketball. How many students play a sport?

*Solution.*

Let  $S$  be the set of students who play soccer and  $B$  be the set of students who play basketball.

Then,  $|S \cup B| = |S| + |B|$  \_\_\_\_\_.



## Principle of Inclusion-Exclusion

When  $A = A_1 \cup \cdots \cup A_k \subset \mathcal{U}$  ( $\mathcal{U}$  for universe) and the sets  $A_i$  are *pairwise disjoint*, we have  $|A| = |A_1| + \cdots + |A_k|$ .

When  $A = A_1 \cup \cdots \cup A_k \subset \mathcal{U}$  and the  $A_i$  are **not** pairwise disjoint, we must apply the **principle of inclusion-exclusion** to determine  $|A|$ :

$$|A_1 \cup A_2| = |A_1| + |A_2| - |A_1 \cap A_2|$$

$$|A_1 \cup A_2 \cup A_3| = |A_1| + |A_2| + |A_3| - |A_1 \cap A_2| - |A_1 \cap A_3| \\ - |A_2 \cap A_3| + |A_1 \cap A_2 \cap A_3|$$

$$|A_1 \cup \cdots \cup A_m| = \sum |A_i| - \sum |A_i \cap A_j| + \sum |A_i \cap A_j \cap A_k| \cdots$$

It may be more convenient to apply inclusion/exclusion where the  $A_i$  are *forbidden* subsets of  $\mathcal{U}$ , in which case \_\_\_\_\_.

## mmm...PIE

The key to using the principle of inclusion-exclusion is determining the right choice of  $A_i$ . The  $A_i$  and their intersections should be easy to count and easy to characterize.

*Notation:*  $\pi = p_1 p_2 \cdots p_n$  is the **one-line notation** for a permutation of  $[n]$  whose first element is  $p_1$ , second element is  $p_2$ , etc.

**Example.** How many permutations  $p = p_1 p_2 \cdots p_n$  are there in which at least one of  $p_1$  and  $p_2$  are even?

*Solution.* Let  $\mathcal{U}$  be the set of  $n$ -permutations.

Let  $A_1$  be the set of permutations where  $p_1$  is even.

Let  $A_2$  be the set of permutations where  $p_2$  is even.

In words,  $A_1 \cap A_2$  is the set of  $n$ -permutations \_\_\_\_\_

Now calculate:  $|A_1| =$   $|A_2| =$

$|A_1 \cap A_2| =$

Applying PIE: So  $|A_1 \cup A_2| =$

## mmm...PIE

**Example.** Find the number of integers between 1 and 1000 that are **not** divisible by 5, 6, or 8.

**Solution.** Let  $\mathcal{U} = \{n \in \mathbb{Z} \text{ such that } 1 \leq n \leq 1000\}$ .

Let  $A_1 \subset \mathcal{U}$  be the multiples of 5,  $A_2 \subset \mathcal{U}$  be the multiples of 6, and  $A_3 \subset \mathcal{U}$  be the multiples of 8. We want  $|\mathcal{U}| - |A_1 \cup A_2 \cup A_3|$ .

In words,  $A_1 \cap A_2$  is the set of integers

$A_1 \cap A_3$  is

$A_2 \cap A_3$  is

and  $A_1 \cap A_2 \cap A_3$  is the set of integers that are

Now calculate:  $|A_1| =$   $|A_2| =$   $|A_3| =$

$|A_1 \cap A_2| =$   $|A_1 \cap A_3| =$   $|A_2 \cap A_3| =$

$|A_1 \cap A_2 \cap A_3| =$

And finally: So  $|\mathcal{U}| - |A_1 \cup A_2 \cup A_3| =$

# Combinations with Repetitions

## Quick review

1. How many ways are there to choose  $k$  elements out of the set  $\{1 \cdot a_1, 1 \cdot a_2, \dots, 1 \cdot a_n\}$ ?
2. How many ways are there to choose  $k$  elements out of the set  $\{k \cdot a_1, k \cdot a_2, \dots, k \cdot a_n\}$ ? (really  $\{\infty \cdot a_1, \infty \cdot a_2, \dots, \infty \cdot a_n\}$ )

What we would like to calculate is:

In how many ways can we choose  $k$  elements out of an arbitrary multiset?

Now, it's as easy as PIE.

## Combinations with Repetitions

**Example.** Determine the number of 10-combinations of the multiset  $S = \{3 \cdot a, 4 \cdot b, 5 \cdot c\}$ .

**Game plan:** Let  $\mathcal{U}$  be the set of 10-combs of  $\{\infty \cdot a, \infty \cdot b, \infty \cdot c\}$ . Use PIE to remove the 10-combs that violate the conditions of  $S$

Define  $A_1$  to be 10-combs that include at least \_\_\_  $a$ 's.

Define  $A_2$  to be 10-combs that include at least \_\_\_  $b$ 's.

Define  $A_3$  to be 10-combs that include at least \_\_\_  $c$ 's.

In words,  $A_1 \cap A_2$  are those 10-combs that

$A_1 \cap A_3$ :

$A_2 \cap A_3$ :

$A_1 \cap A_2 \cap A_3$

Now calculate:  $|\mathcal{U}| = \quad |A_1| = \quad |A_2| = \binom{3}{5} \quad |A_3| = \binom{3}{4}$   
 $|A_1 \cap A_2| = 3 \quad |A_1 \cap A_3| = 1 \quad |A_2 \cap A_3| = 0 \quad |A_1 \cap A_2 \cap A_3| = 0$

And finally: So  $|\mathcal{U}| - |A_1 \cup A_2 \cup A_3| =$

# Derangements

At a party, 10 partygoers check their hats. They “have a good time”, and are each handed a hat on the way out. In how many ways can the hats be returned so that no one is returned his/her own hat?

This is a **derangement** of ten objects.

*Definition:* An  **$n$ -derangement** is an  $n$ -permutation  $\pi = p_1 p_2 \cdots p_n$  such that  $p_1 \neq 1, p_2 \neq 2, \dots, p_n \neq n$ .

*Note:* A derangement is a permutation without fixed points  $\pi(i) = i$ .

*Notation:* We let  $D_n$  be the number of all  $n$ -derangements.

When you see  $D_n$ , think combinatorially: “The number of ways to return  $n$  hats to  $n$  people so no one gets his/her own hat back”

# Calculating the number of derangements

**Example.** Calculate  $D_n$ .

**Solution.** Let  $\mathcal{U}$  be the set of all  $n$ -permutations.

Remove bad permutations using PIE.

For all  $i$  from 1 to  $n$ , define  $A_i$  to be  $n$ -perms where  $p_i = i$ .

In words,  $A_i \cap A_j$  are  $n$ -perms where

$A_i \cap A_j \cap A_k$  are  $n$ -perms where

In general,  $A_{i_1} \cap \cdots \cap A_{i_k}$  are  $n$ -perms with  $p_{i_1} = i_1, \cdots, p_{i_k} = i_k$ .

Now calculate:  $|\mathcal{U}| =$   $|A_1| =$   $|A_2| =$

**For all**  $i$  and  $j$ ,  $|A_i \cap A_j| =$

When intersecting  $k$  sets,  $|A_{i_1} \cap \cdots \cap A_{i_k}| =$

Recall:  $|A_1 \cup \cdots \cup A_n| = \sum |A_i| - \sum |A_i \cap A_j| + \sum |A_i \cap A_j \cap A_k| \cdots$

Therefore,  $D_n = |\mathcal{U}| - |A_1 \cup \cdots \cup A_n| =$



## Randomly returning hats

Upon simplification, we see

$$\begin{aligned}
 D_n &= n! - \binom{n}{1}(n-1)! + \binom{n}{2}(n-2)! - \cdots + (-1)^n \binom{n}{n} 0! \\
 &= n! - \frac{n!}{1!} + \frac{n!}{2!} - \cdots + (-1)^n \frac{n!}{n!} \\
 &= n! \left[ 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^n \frac{1}{n!} \right]
 \end{aligned}$$

**Recall:** Taylor series expansion of  $e^x$ :

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

Plug in  $x = -1$  and truncate after  $n$  terms to see that

$$e^{-1} \approx \left[ 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^n \frac{1}{n!} \right]$$

**Conclusion:** If  $n$  people go to a party and the hats are passed back randomly, the probability that no one gets his or her hat back at the party is  $D_n/n!$ , which is approximately  $1/e \approx 37\%$ .

## Combinatorial proof involving $D_n$

**Recall:** The combinatorial interpretation of  $D_n$  is: “The number of ways to return  $n$  hats to  $n$  people so no one gets his/her own hat back”

**Example.** Prove the following recurrence relation for  $D_n$  combinatorially.

$$D_n = (n - 1)(D_{n-2} + D_{n-1})$$

## A formula for Stirling numbers (p. 90)

(Careful: change of notation !!)

Recall:  $S(n, k) = \left\{ \begin{matrix} n \\ k \end{matrix} \right\}$  is the number of **set partitions** of  $[n]$  into exactly  $k$  parts, and  $k!S(n, k)$  is the number of **onto functions**  $[n] \rightarrow [k]$ .

*Question:* What is a formula for  $S(n, k)$ ?

*Solution.* We will find the number of surjections from  $[n]$  to  $[k]$ .

Use PIE with  $\mathcal{U}$  = set of **all** functions from  $[n]$  to  $[k]$ .

We will remove the “bad” functions where the range is not  $[k]$ .

Define  $A_i$  be the set of functions  $f : [n] \rightarrow [k]$  where  $i$  is not “hit”.

In words,  $A_{i_1} \cap \cdots \cap A_{i_j}$  are functions where none of  $i_1$  through  $i_j$  are elements of the image.

We calculate:  $|\mathcal{U}| = k^n$ ,  $|A_i| = (k - 1)^n$ ,  $|A_i \cap A_j| = (k - 2)^n$

When intersecting  $j$  sets,  $|A_{i_1} \cap \cdots \cap A_{i_j}| = (k - j)^n$ .

Therefore,  $k!S(n, k) = \sum_{j=0}^k (-1)^j \binom{k}{j} (k - j)^n$ ; we conclude

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^j \binom{k}{j} (k - j)^n = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

# A formula for Bell numbers (p. 166)

(Careful: change of notation !!)

Recall:  $B_n$  is the number of partitions of  $[n]$  into any number of parts. Manipulate our expression from prev. page to find a formula.

$$\begin{aligned}
 B_n &= \sum_{k \geq 0} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \sum_{k \geq 0} \frac{1}{k!} \sum_{j=0}^k \frac{k!}{j!(k-j)!} (-1)^{k-j} j^n \\
 &= \sum_{k \geq 0} \sum_{j=0}^k \frac{1}{j!(k-j)!} (-1)^{k-j} j^n = \sum_{k \geq 0} \sum_{j=0}^k \frac{(-1)^{k-j} j^n}{(k-j)! j!} \\
 &= \sum_{j \geq 0} \sum_{k \geq j} \frac{(-1)^{k-j} j^n}{(k-j)! j!} = \sum_{j \geq 0} \frac{j^n}{j!} \sum_{m \geq 0} \frac{(-1)^m}{(m)!} = \sum_{j \geq 0} \frac{j^n}{j!} \frac{1}{e}
 \end{aligned}$$

**Theorem 4.3.3.** For any  $n > 0$ ,  $B_n = \frac{1}{e} \sum_{j \geq 0} \frac{j^n}{j!}$ .

For example,  $B_5 = \frac{1}{e} \left( \frac{0^5}{0!} + \frac{1^5}{1!} + \frac{2^5}{2!} + \frac{3^5}{3!} + \frac{4^5}{4!} + \frac{5^5}{5!} + \dots \right) = 52$ .